



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> C07H 21/00, C12N 15/00 A23C 9/00, A01N 1/02	<b>A1</b>	<b>(11) International Publication Number:</b> WO 93/25567 <b>(43) International Publication Date:</b> 23 December 1993 (23.12.93)
<b>(21) International Application Number:</b> PCT/US93/05724 <b>(22) International Filing Date:</b> 15 June 1993 (15.06.93) <b>(30) Priority data:</b> 07/898,956 15 June 1992 (15.06.92) US <b>(60) Parent Application or Grant</b> (63) Related by Continuation US 07/898,956 (CIP) Filed on 15 June 1992 (15.06.92) <b>(71) Applicant (for all designated States except US):</b> GEN- PHARM INTERNATIONAL, INC. [US/US]; 297 North Bernardo Avenue, Mountain View, CA 94043 (US).		<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DEBOER, Herman, A. [NL/NL]; Sotaweg 24, NL-2371 GD Roelofanendsveen (NL). STRIJKER, Rein [NL/NL]; Spaargarenstaat 2, NL-2341 JW Oegstgeest (NL). HEYNEKER, Herbert, L. [US/US]; 501 Roehampton Road, Hillsborough, CA 94010 (US). PLATENBURG, Gerard [NL/NL]; Wijn- gaardenlaan 56, NL-2252 XR Voorschoten (NL). LEE, Sang, He [NL/NL]; Fien de la Marstraat 26, NL-2331 HN Leiden (NL). PIEPER, Frank [NL/NL]; C. van Maasdijkstraat 13, NL-3555 VM Utrecht (NL). <b>(74) Agent:</b> SMITH, William, M.; Townsend and Townsend Khourie and Crew, One Market Plaza, 20th Fl., Steuart Tower, San Francisco, CA 94105 (US). <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PRODUCTION OF RECOMBINANT POLYPEPTIDES BY BOVINE SPECIES AND TRANSGENIC METH- ODS		
<b>(57) Abstract</b> <p>A transgene for producing recombinant polypeptides in the milk of transgenic bovine species comprises at least one ex-          pression regulation sequence, a secretory DNA sequence encoding a secretory signal sequence which is functional in mammary          secretory cells of the bovine species and a recombinant DNA sequence encoding the recombinant polypeptide. Also included are          methods for producing transgenic bovine species. The method includes introducing the above transgene into an embryonal target          cell of a bovine species, transplanting the transgenic embryonic target cell formed thereby into a recipient bovine parent and in-          dentifying at least one female offspring which is capable of producing the recombinant polypeptide in its milk. The invention al-          so includes transgenic bovine species as well as the milk from such transgenic bovine species. Methods are also provided for          producing transgenic non-human mammals having a desirable phenotype. The method comprises first methylating a transgene          followed by introduction into fertilized oocytes. The oocytes are then cultured to form pre-implantation embryos. Thereafter, at          least one cell is removed from each of the pre-implantation embryos and the DNA digested with a restriction endonuclease cap-          able of cleaving the methylated transgene but incapable of cleaving the unmethylated form of the transgene.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

PRODUCTION OF RECOMBINANT POLYPEPTIDES  
BY BOVINE SPECIES AND TRANSGENIC METHODS

5

Field of the Invention

The invention relates to the production of recombinant polypeptides by transgenic bovine species and to methods  
10 for producing transgenic non-human mammals having a desired phenotype.

Background of the Invention

There is a plethora of literature relating to the  
15 expression of heterologous genes in lower organisms such as unicellular bacteria, yeast and filamentous fungi, and in higher cell types such as mammalian cells. There are also numerous reports on the production of transgenic animals, most of which relate to the  
20 production of transgenic mice. See, e.g., U.S. Pat. No. 4,736,866 (transgenic mice containing activated oncogene); Andres, A., et al. (1987) Proc. Natl. Acad. Sci. USA 84:1299-1303 (HA-RAS oncogene under control of whey acid protein promoter); Schoenberger, C.A., et al.

-2-

(1987) Experientia 43:644 and (1988) EMBO J. 7:169-175 (C-myc oncogene under control of whey acid protein promoter); and Muller, W.J., et al. (1988) Cell 54:105-115 (C-myc oncogene under control of the mouse mammary tumor virus promoter). Several laboratories have also reported the production of transgenic porcine species (Miller, K.F., et al. (1989) J. Endocrin. 120:481-488 (expression of human or bovine growth hormone gene in transgenic swine); Vize, P.D., et al. (1988) J. Cell Sci. 90:295-300 (porcine growth hormone fusion gene in transgenic pigs); and Ebert, K. et al. (1988) Mol. Endocrin. 2:277-283 (MMLV-rat somatotropin fusion gene in transgenic pigs)), transgenic sheep (Nancarrow, et al. (1987) Theriogenology 27:263 (transgenic sheep containing bovine growth hormone gene) Clark, A.J. et al. (1989) Bio/Technology 7:487-482 and Simons, J., et al. (1988) Bio/Technology 6:179-183 (human factor IX and  $\alpha$ -1 antitrypsin CONA in ovine species), and rabbit (Hanover, S.V., et al. (1987) Deutsche Tierärztliche Wochenschrift 94,:476-478 (production of transgenic rabbits by injection of uteroglobin-promoter-CAT fusion gene into fertilized rabbit oocytes). A number of reports have also suggested the production of transgenic cattle (Wagner, et al. (1984) Theriogenology 21:29-44) with one reporting some progress in microinjection techniques (Lohse, J.K., et al. (1985) Theriogenology 23:205). However, little, if any, success has been achieved in producing transgenic cows. Scientific articles which clearly demonstrate the actual production of a transgenic cow capable of producing a heterologous protein are presently unknown. This, despite the statements that one transgenic cow was produced in Canada which expressed human  $\beta$ -interferon (Van Brunt, J. (1988) Bio/Technology 6:1149-1155) and that transient expression of human  $\alpha$ -fetoprotein in liver and blood was obtained on one occasion (Church, R.B. (1986) Biotechnology News Watch 6 (15), 4). One reference

-3-

reports that bovine papilloma virus was apparently integrated but not expressed in a transgenic cow (Roschlau, et al. (1988) Arch. Tierz., Berlin 31:3-8). A recent article has summarized the genetic engineering of livestock. (Pursel, V.G. et al. (1989) Science 244:1281-1288).

A number of laboratories have reported tissue-specific expression of DNA encoding various proteins in the mammary gland or the production of various proteins in the milk of transgenic mice and sheep. For example, Simmons, J.P., et al. (1987) Nature 328:530-532 report the microinjection of a 16.2 kb genomic fragment encoding  $\beta$ -lactoglobulin (BLG) including 4 kb of 5' sequence, 4.9 kb of the BLG transcription unit and 7.3 kb of 3' flanking sequence into fertilized mouse eggs. According to these authors, the sheep BLG was expressed in mammary tissue and produced BLG in the milk of the transgenic mice at concentrations ranging from about 3.0 to about 23 mg/ml. When, however, cDNA encoding human factor IX or human  $\alpha$ 1-antitrypsin was inserted into the 5' untranslated region of the BLG gene and microinjected into sheep (Simmons, J.P., et al. (1988) Bio/Technology 6:179-183) the production of factor IX or  $\alpha$ 1-antitrypsin was significantly reduced (25ng/ml for factor IX and 10mg/ml for  $\alpha$ 1-antitrypsin; see Clark, A.J., et al. (1989) Bio/Technology 7:487-492).

In a similar approach, a 14 kb genomic clone containing the entire 7.5 kb rat  $\beta$ -casein together with 3.5 kb of 5' and 3.0 kb of 3' flanking DNA was reportedly microinjected into fertilized mouse oocytes. Lee, et al. (1988) Nucl. Acids Res. 16:1027-1041. Yet, in this case, the level of expression of the rat  $\beta$ -transgene in the lactating mammary gland of transgenic mice was

-4-

reported to be at a level of 0.01-1% of the endogenous mouse  $\beta$ -casein gene.

Human tissue plasminogen activator (t-PA) reportedly was produced in transgenic mouse milk at the levels between  
5 0.2 and about 0.4  $\mu$ g/ml when a cDNA encoding a human t-PA with its endogenous secretion sequence was expressed under control of a 2.6 kb 5' sequence of the murine whey acid protein gene. Gordon, K., et al. (1987) Bio/Technology 5:1183-1187. Subsequent experiments  
10 using the same or similar construction reportedly produced t-PA in different mouse lines arranging from less than 20ng of t-PA per ml of milk to about 50  $\mu$ g/ml. Pittius, C.W., et al. (1988) Proc. Natl. Acad. Sci. USA 85:5874-5878.

15 U.S. Patent No. 4,873,316 issued October 10, 1989, discloses the use of 9 kb of 5' sequence from the bovine  $\alpha$ S1 casein gene including the casein signal peptide and several casein codons fused to a mature t-PA sequence. The transgenic mice obtained with this construct  
20 reportedly produced about 0.2-0.5  $\mu$ g/ml of a t-PA fusion protein in their milk.

In addition, a number of patent publications purportedly describe the production of specific proteins in the milk of transgenic mice and sheep. See, e.g. European Patent  
25 Publication No. 0 264 166 published April 20, 1988 (hepatitis B surface antigen and t-PA genes under control of the whey acid promoter protein for mammary tissue specific expression in mice); PCT Publication No. W088/00239 published January 14, 1988 (tissue specific  
30 expression of a transgene encoding factor IX under control of a whey protein promoter in sheep); PCT Publication No. W088/01648 published March 10, 1988 (transgenic mouse having mammary secretory cells incorporating a recombinant expression system comprising

-5-

a bovine  $\alpha$ -lactalbumin gene fused to interleukin-2); European Pat. Pub. No. 0 279 582 published August 24, 1988 (tissue-specific expression of chloramphenicol acetyltransferase under control of rat  $\beta$ -casein promoter  
5 in transgenic mice); and PCT Pub. No. W088/10118 published December 29, 1988 (transgenic mice and sheep containing transgene encoding bovine  $\alpha$ S1 casein promoter and signal sequence fused to t-PA).

10 Given the state of the transgenic art, it is apparent that a need exists for methods which enable the efficient production of transgenic mammals, especially transgenic mammals other than transgenic mice.

15 Further, it is apparent that a need exists for methods for producing transgenic bovine species which are capable of producing recombinant polypeptides such as human milk proteins and human serum proteins in the milk of such transgenic mammals.

20 Accordingly, it is an object herein to provide methods for detecting the transgenesis of fertilized oocytes prior to implantation.

In addition, it is an object herein to provide transgenic bovine species which are capable of producing recombinant polypeptides which are maintained intracellularly or are secreted extracellularly.

25 It is also an object herein to provide transgenic bovine species which are capable of producing recombinant polypeptides such as human milk proteins and human serum proteins in the milk of such transgenic animals.

30 Further, it is an object herein to provide milk from a transgenic bovine species containing such recombinant polypeptides.

-6-

Still further, it is an object herein to provide food formulations supplemented with recombinant polypeptides from such transgenic milk such as human infant formula supplemented with human lactoferrin.

- 5 Further, it is an object herein to provide transgenes which are capable of directing the production of recombinant polypeptides in the milk of transgenic bovine species.

10 The references discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the inventors are not entitled to antedate such disclosure by priority based on earlier filed applications.

15 Summary of the Invention

In accordance with the above objects, the invention includes transgenes for producing recombinant polypeptides in the milk of transgenic bovine species. The production of such transgenic bovine milk containing  
20 one or more recombinant polypeptides is desirable since it provides a matrix wherein little or no purification is necessary for human consumption. The transgene comprises a secretory DNA sequence encoding a secretory signal sequence which is functional in mammary secretory  
25 cells of the bovine species of interest and a recombinant DNA sequence encoding the recombinant polypeptide. These sequences are operably linked to form a secretory-recombinant DNA sequence. At least one expression regulation sequence, functional in the  
30 mammary secretory cells of the bovine species, is operably linked to the secretory-recombinant DNA sequence. The transgene so constructed is capable of directing the expression of the secretory-recombinant DNA sequence in mammary secretory cells of bovine



-7-

species containing the transgene. Such expression produces a form of recombinant polypeptide which is secreted from the mammary secretory cells into the milk of the transgenic bovine species.

- 5 In addition, the invention includes methods for producing such transgenic bovine species. The method includes introducing the above transgene into an embryonal target cell of a bovine species, transplanting the transgenic embryonic target cell formed thereby into  
10 a recipient bovine parent and identifying at least one female offspring which is capable of producing the recombinant polypeptide in its milk.

The invention also includes transgenic bovine species capable of producing recombinant polypeptides in the  
15 milk of lactating females of said species, the milk from such transgenic bovine species containing such recombinant polypeptides and food formulations containing the transgenic milk in liquid or dried form, as well as food formulations supplemented with one or  
20 more recombinant polypeptides from such transgenic milk.

In addition to the foregoing, the invention includes transgenes and transgenic bovine species containing transgenes that are capable of producing a recombinant polypeptide. Such transgenes are similar to the  
25 aforementioned transgenes for milk secretion and are characterized by having an expression regulation sequence which targets the expression of the DNA encoding the recombinant polypeptide to a particular cell or tissue type, e.g. expression of human serum  
30 albumin in the liver of a transgenic bovine species. When the recombinant polypeptide is to be secreted from such targeted cells or tissues, a secretory DNA sequence encoding a secretory signal sequence functional in the

-8-

particular targeted cell or tissue is operably linked to the recombinant DNA sequence encoding the recombinant polypeptide, e.g. secretion of human serum albumin from bovine liver into the bovine circulatory system.

- 5 Further, the invention includes methods for producing transgenic non-human mammals having a desirable phenotype. The method comprises first causing the methylation of a transgene capable of conferring the desirable phenotype when incorporated into the cells of
- 10 a transgenic non-human animal, e.g., by transforming an appropriate bacterium, such as E. coli MM 294, with a plasmid containing the transgene. The methylated transgene is then excised and introduced into fertilized oocytes of the non-human animal to permit integration
- 15 into the genome. The oocytes are then cultured to form pre-implantation embryos thereby replicating the genome of each of the fertilized oocytes. Thereafter, at least one cell is removed from each of the pre-implantation embryos and treated to release the DNA contained
- 20 therein. Each of the released DNAs are then digested with a restriction endonuclease capable of cleaving the methylated transgene but incapable of cleaving the unmethylated form of the transgene formed after integration into and replication of the genomic DNA.
- 25 Those pre-implantation embryos which have integrated the transgene contain DNA which is resistant to cleavage by the restriction endonuclease in the region containing the transgene. This resistance to digestion, which can be detected by electrophoresis of the digest after PCR
- 30 amplification of the DNA and hybridization with a labelled probe for the transgene, facilitates the identification of successful transgenesis.

The invention also includes a method to produce a population of transgenic offspring having the same

35 genotype. This method utilizes a specific embodiment

-9-

of the above method for detecting early transgenesis. In this method, a methylated transgene is introduced into fertilized oocytes which are cultured to pre-implantation embryos. Thereafter, each pre-implantation embryo is divided to form first and second hemi-embryos. Each of the first hemi-embryos are then analyzed for transgenesis as described above. After identifying successful transgenesis in at least one first hemi-embryo, the second untreated hemi-embryo which contains the integrated transgene, is cloned to form a multiplicity of clonal transgenic blastocysts or hemi-blastocysts, each of which have the same genotype. The transgenic embryos are thereafter transplanted into one or more recipient female parents to produce a population of transgenic non-human mammals having the same genotype.

#### Brief Description of the Drawings

The accompanying drawings, which are incorporated in and form a part of the specification, illustrate embodiments of the present invention and, together with the description, serve to explain the principles of the invention. In the drawings:

Fig. 1 depicts the DNA (Seq. ID No.: 1) and amino acid (Seq. ID No.: 2) sequence for a human lactoferrin clone derived from a human mammary cDNA library as described herein except that the sequence between nucleotides 1557-1791 and 2050-2119 corresponds to the previously published sequence (Rado et al. (1987) Blood 70:989-993).

Fig. 2 depicts the complete DNA (Seq. ID No.: 3) and amino acid (Seq. ID No.: 4) sequence of human lactoferrin including 5' and 3' untranslated sequence as well as the complete human lactoferrin signal sequence.

-10-

Fig. 3 is a restriction map of a clone of a 5'-flanking region of bovine  $\alpha$ S1 casein gene.

Fig. 4 is a restriction map of a clone of a 3'-flanking region of bovine  $\alpha$ S1 casein gene.

5 Figs. 5A, 5B and 5C depict the construction of pSI3'5'CAT and pSI5'CAT.

Fig. 6 depicts pMH-1.

Figs. 7A through 7F depict the construction of expression vectors containing sequences encoding human  
10 lactoferrin.

Fig. 8 depicts the genome of human serum albumin, the fragments used to generate transgenic mice contained in this genomic DNA and the identification of the fragment sizes which would be obtained upon the digestion of  
15 genomic DNA from a transgenic mouse with the restriction enzymes BstE-II and Nco-I or with Nco-I and Hindi-III.

Fig. 9 depicts an alternate pathway for the construction of a transgene of the invention encoding human lactoferrin.

20 Fig. 10 depicts the construction of a plasmid pPC containing a transgene encoding Protein C.

Fig. 11 depicts the DNA sequence for a hybrid intervening sequence used in a preferred embodiment of the invention. This hybrid sequence comprises a 5' portion from an intervening sequence of bovine  $\alpha$ S1 casein and a 3' portion from an intervening sequence of an IgG intervening sequence. The juncture of the 5' and  
25 3' portion is the HindIII site shown.

-11-

Fig. 12A is a restriction map of a bovine  $\alpha$ S1 casein promoter hLF cDNA transgene.

Fig. 12B shows a Southern blot analysis of DNA isolated from various bovine and murine tissues using an hLF cDNA  
5 probe.

Fig. 13 depicts restriction maps of hLF genomic clones 13.1 and 13.2.

Fig. 14 depicts the BamHI fragment from genomic hLF subcloned into plasmid pUC19.

10 Fig. 15A depicts a restriction map of the 8hLFgen9k or 16hLFgen9k construct containing the 8 or 16 kb  $\alpha$ S1 casein promoter, a ClaI-ApaI synthetic linker and the 9 kb (i.e., 8.9 kb) ApaI-SalI genomic hLF fragment.

Fig. 15B depicts the DNA sequence of the ClaI-ApaI  
15 synthetic sequence shown in Fig. 15A.

Fig. 15C depicts the IVS and the structure of exon 1 and part of exon 2 of the genomic hLF construct shown in Fig. 15A through Fig. 17.

Fig. 16 depicts the coinjection of the NotI-SalI  
20 fragment from the 8hLFgen9k or 16hLFgen9k construct (as shown in Fig. 15A) with the 3' ClaI fragment of genomic hLF.

Fig. 17 depicts the generation of a genomic 8hLF transgene by linking the NotI-MluI fragment from the  
25 8hLFgen9k construction (shown in Fig. 15A), the MluI-ClaI fragment from clone 13.2 depicted in Fig. 13 and a ClaI-NotI linker. Fig. 17 also depicts the DNA sequence of the ClaI-NotI linker.

-12-

Figs. 18-20 depict the generation of the  $\beta$ LG-hLFgen and  $\beta$ LG-hLFgen37 constructs.

Fig. 21 depicts the design of the 16,8hLZ expression vector.

5 Fig. 22 depicts the design of the 16,8hLZ3 expression vector.

Fig. 23A-23E depict the pathway for the construction of plasmid p16,8hLZ.

10 Fig. 24 depicts a comparison between the DNA of bovine  $\beta$ LG and sheep  $\beta$ LG. The top sequence represents the bovine sequence.

Fig. 25 shows the linker GP 278/279.

Fig. 26 depicts the p16,8A hLZ3 expression vector.

Fig. 27 depicts the 16,A hLZ3 expression vector.

15 Detailed Description of the Invention

The "non-human mammals" of the invention comprise all non-human mammals capable of producing a "transgenic non-human mammal" having a "desirable phenotype". Such mammals include non-human primates, murine species,  
20 bovine species, canine species, etc. Preferred non-human animals include bovine, porcine and ovine species, most preferably bovine species.

Desirable phenotypes for transgenic non-human mammals include, but are not limited to, the production of  
25 recombinant polypeptides in the milk of female transgenic non-human mammals, the production of animal models for the study of disease, the production of

-13-

animals with higher resistance to disease (e.g. diseases of the mammary gland such as mastitis) and the production of recombinant polypeptides in the blood, urine or other suitable body fluid or tissue of the animal. In the preferred embodiments, transgenic bovine species are disclosed which are capable of producing recombinant human lactoferrin, human serum albumin and human Protein C in the milk of lactating females or human serum albumin in the liver of the transgenic animal.

The transgenic non-human mammals of the invention are produced by introducing a "transgene" into an embryonal target cell of the animal of choice. In one aspect of the invention, a transgene is a DNA sequence which is capable of producing a desirable phenotype when contained in the genome of cells of a transgenic non-human mammal. In specific embodiments, the transgene comprises a "recombinant DNA sequence" encoding a "recombinant polypeptide". In such cases, the transgene is capable of being expressed to produce the recombinant polypeptide.

As used herein, a "recombinant polypeptide" (or the recombinant DNA sequence encoding the same) is either a "heterologous polypeptide" or a "homologous polypeptide". Heterologous polypeptides are polypeptides which are not normally produced by the transgenic animal. Examples of heterologous polypeptides include human milk proteins such as lactoferrin, lysozyme, secreted immunoglobulins, lactalbumin, bile salt-stimulated lipase, etc., human serum proteins such as albumin, immunoglobulins, Factor VIII, Factor IX, protein C, etc. and industrial enzymes such as proteases, lipases, chitinases, and ligninases from procaryotic and eucaryotic sources. The

-14-

recombinant DNA sequences include genomic and cDNA sequences encoding the recombinant polypeptide.

When recombinant DNA sequences encoding a heterologous polypeptide are used, the transgene may be integrated  
5 in a random manner into the genome of the species used for transgenesis. As disclosed in the Examples, transgenes encoding human lactoferrin, human serum albumin and human Protein C in conjunction with a  $\alpha$ S1 casein secretory signal sequence under control of  $\alpha$ S1  
10 casein expression regulation sequences are designed to produce and secrete these heterologous polypeptides from the mammary gland of a lactating transgenic mammal into its milk.

As used herein, a homologous polypeptide is one which  
15 is endogenous to the particular transgenic species. Examples of endogenous polypeptides from bovine species include bovine milk proteins such as  $\alpha$ S1,  $\alpha$ S2,  $\beta$ - and  $\kappa$ -casein,  $\beta$ -lactoglobulin lactoferrin, lysozyme, cholesterol hydrolase, serum proteins such as serum  
20 albumin and proteinaceous hormones such as growth hormones. When recombinant DNA sequences encoding a homologous polypeptide are used, the transgene is preferably integrated in a random manner into the genome of the species used for transgenesis. Such random  
25 integration results in a transgenic animal which contains not only the transgene encoding the endogenous polypeptide but also the corresponding endogenous genomic DNA sequence. Accordingly, such transgenic non-human mammals are readily characterized by an increase  
30 in the copy number of genes encoding the endogenous polypeptide. Further, the transgene will generally be located at a position which is different from the endogenous gene.



-15-

When DNA encoding a homologous polypeptide is expressed, for example, in bovine species, the transgenic animal is characterized by an increase in the amount of the homologous polypeptide in either the endogenous tissue or fluid in which it is normally found and/or by its presence in a tissue and/or body fluid which either does not normally contain the homologous polypeptide or produces it at significantly lower levels.

Thus, for example, bovine cholesterol hydrolase is normally present in the colostrum for about the first 15-20 days of lactation. This naturally occurring endogenous polypeptide increases calf weight. This protein, however, is also a homologous polypeptide when, for example, its expression in mammary secretory cells is placed under the control of expression regulation sequences, such as those obtained from bovine casein genes, which facilitate the expression of the homologous polypeptide beyond the lactation period that it is normally present. Thus, according to one aspect of the invention, bovine cholesterol hydrolase expression is maintained in transgenic bovine milk by placing the expression of cholesterol hydrolase recombinant DNA (either cDNA or genomic) under the control of bovine  $\alpha$ S1 casein expression regulation sequences. When a genomic recombinant DNA is used, it is engineered such that it has appropriate restriction sites (e.g. ClaI and SalI) at the 5' and 3' end of the structural gene such that it is capable of being inserted into an appropriate transgene genomic cassette (e.g. p-16 kb, CS which is described in Example 15). Alternatively, a recombinant DNA encoding bovine cholesterol hydrolase derived from cDNA may be placed under control of bovine  $\alpha$ S1 casein expression regulation sequence by substituting the human lactoferrin sequences in a plasmid such as p16, 8HLF3 (containing a hybrid intervening sequence) or p16, 8HLF4 (containing a homologous  $\alpha$ S1 casein intervening

-16-

sequence). When these particular plasmids are used, the cDNA clone is engineered such that it has appropriate ClaI and SalI restriction sites at the ends of the recombinant DNA.

- 5 By way of further example, bovine lactoferrin is normally present in only trace amounts in cow's milk. When, however, bovine lactoferrin is expressed under control of other regulatory sequences, for example, obtained from an  $\alpha$ S1 casein gene, higher amounts of
- 10 lactoferrin in the milk of transgenic bovine species are obtained. In another example, a transgene comprising DNA encoding homologous bovine growth hormone is incorporated into the bovine genome to confer superior growth characteristics to the transgenic animal. In
- 15 other instances, homologous polypeptides include, for example, a polypeptide which normally is maintained intracellularly in a particular species but which is secreted into the milk or other extracellular compartment of the transgenic species, such as the
- 20 circulatory system.

- Each of the heterologous or homologous polypeptides are characterized by specific amino acid and nucleic acid sequences. It is to be understood, however, that such sequences include naturally occurring allelic variations
- 25 thereof and variants produced by recombinant methods wherein such nucleic acid and polypeptide sequences have been modified by the substitution, insertion and/or deletion of one or more nucleotides in such nucleic acids to cause the substitution, insertion or deletion
- 30 of one ore more amino acid residues in the recombinant polypeptide.

When expression of the DNA of the transgene is necessary to generate a desired phenotype, e.g. to produce a recombinant polypeptide, the transgene typically

-17-

includes at least a 5' and preferably additional 3' "expression regulation sequences" each operably linked to a recombinant or secretory-recombinant DNA as defined hereinafter. Such expression regulation sequences in addition to controlling transcription also contribute to RNA stability and processing, at least to the extent they are also transcribed.

Such expression regulation sequences are chosen to produce tissue-specific or cell type-specific expression of the recombinant or secretory-recombinant DNA. Once a tissue or cell type is chosen for expression, 5' and optional 3' expression regulation sequences are chosen. Generally, such expression regulation sequences are derived from genes that are expressed primarily in the tissue or cell type chosen. Preferably, the genes from which these expression regulation sequences are obtained are expressed substantially only in the tissue or cell type chosen, although secondary expression in other tissue and/or cell types is acceptable if expression of the recombinant DNA in the transgene in such tissue or cell type is not detrimental to the transgenic animal. Particularly preferred expression regulation sequences are those endogenous to the species of animal to be manipulated. However, expression regulation sequences from other species such as those from human genes may also be used. Particularly preferred expression regulation sequences from human genes are human lactoferrin (hLF) sequences. In some instances, the expression regulation sequences and the recombinant DNA sequences (either genomic or cDNA) are from the same species, e.g., each from bovine species or from a human source. In such cases, the expression regulation sequence and the recombinant DNA sequence are homologous to each other. Alternatively, the expression regulation sequences and recombinant DNA sequences (either cDNA or genomic) are obtained from different species, e.g., an

-18-

expression regulation sequence from bovine species and a recombinant DNA sequence from a human source). In such cases, the expression regulation and recombinant DNA sequence are heterologous to each other. The following defines expression regulation sequences from endogenous genes. Such definitions are also applicable to expression regulation sequences from non-endogenous, heterologous genes.

In general, the 5' expression regulation sequence includes the transcribed portion of the endogenous gene upstream from the translation initiation sequence (the 5' untranslated region or 5' UTR) and those flanking sequences upstream therefrom which comprise a functional promoter. As used herein, a "functional promoter" includes those necessary untranscribed DNA sequences which direct the binding of RNA polymerase to the endogenous gene to promote transcription. Such sequences typically comprise a TATA sequence or box located generally about 25 to 30 nucleotides from the transcription initiation site. The TATA box is also sometimes referred to the proximal signal. In many instances, the promoter further comprises one or more distal signals located upstream from the proximal signal (TATA box) which are necessary to initiate transcription. Such promoter sequences are generally contained within the first 100 to 200 nucleotides located upstream from the transcription initiation site, but may extend up to 500 to 600 nucleotides from the transcription initiation site. Such sequences are either readily apparent to those skilled in the art or readily identifiable by standard methods. Such promoter sequences alone or in combination with the 5' untranslated region are referred to herein as "proximal 5' expression regulation sequences".

-19-

In addition to such proximal 5' expression regulation sequences, it is preferred that additional 5' flanking sequences (referred to herein as "distal 5' expression regulation sequences") also be included in the transgene. Such distal 5' expression regulation sequences are believed to contain one or more enhancer and/or other sequences which facilitate expression of the endogenous gene and as a consequence facilitate the expression of the recombinant or secretory-recombinant DNA sequence operably linked to the distal and proximal 5' expression regulation sequences. The amount of distal 5' expression regulation sequence depends upon the endogenous gene from which the expression regulation sequences are derived. In general, however, such sequences comprise 5' flanking regions of approximately 1 kb, more preferably 16 kb and most preferably about 30 kb of 5' flanking sequence. The determination of the optimal amount of distal 5' expression regulation sequence used from any particular endogenous gene is readily determined by varying the amount of distal 5' expression regulation sequence to obtain maximal expression. In general, the distal 5' expression regulation sequence will not be so large as to extend into an adjacent gene and will not include DNA sequences which adversely effect the level of transgene expression.

In addition, it is preferred that 3' expression regulation sequences also be included to supplement tissue or cell-type specific expression. Such 3' expression regulation sequences include 3' proximal and 3' distal expression regulation sequences from an appropriate endogenous gene. The 3' proximal expression regulation sequences include transcribed but untranslated DNA positioned downstream from the translation stop signal in the recombinant DNA sequence (also referred to as the 3' untranslated region or 3'

-20-

UTR). Such sequences generally terminate at a polyadenylation sequence (either from the endogenous gene or from other sources such as SV40) and sequences that may affect RNA stability. Generally, 3' UTR's  
5 comprise about 100 to 500 nucleotides downstream from the translation stop signal in the gene from which the 3' regulation sequence is derived. Distal 3' expression regulation sequences include flanking DNA sequences downstream from the proximal 3' expression regulation  
10 sequence. Some of these distal sequences are transcribed, but do not form part of the mRNA while other sequences in this distal 3' expression regulation sequence are not transcribed at all. Such distal 3' expression regulation sequences are believed to contain  
15 enhancer and/or other sequences which enhance expression. Such sequences are believed to be necessary for efficient polydenylation and contain transcription termination sequences. Preferably, such sequences comprise about 2 kb, more preferably 8 kb and most  
20 preferably about 15 kb of 3' flanking sequence.

A preferred 3' flanking sequence is the 3' flanking sequence of the human lactoferrin (hLF) gene. Transgenic animals containing transgenes that include about 9 kb of hLF 3' flanking sequences show enhanced expression  
25 of recombinant polypeptides in milk compared to animals containing transgenes that include 1 kb or less of hLF 3' flanking sequence, due to an enhancer or other enhancing sequence located in this region. Usually the human lactoferrin 3' flanking sequence will be at least  
30 1 kb in length up to about 9 kb in length or longer, typically 3 to 7 kb, more typically 4 to 5 kb. It will also be possible, and sometimes desirable, to use standard methods (e.g., deletion analysis) to identify regions contained within the 9 kb 3' flanking sequence  
35 that enhance mammary gland expression of recombinant polypeptides. These enhancers or enhancing sequences can

-21-

be isolated and used in combination with various amounts of homologous or heterologous sequences. Typically the enhancing sequences can range in length from about 50 basepairs to about 2 kb, more typically from about 100  
5 basepairs to about 500 basepairs.

It will often be desirable to use a transgene having a 5' expression regulation sequence and a 3' flanking sequence that originate from the same gene. In a preferred embodiment, the 5' expression regulation  
10 sequence and 3' flanking sequence are from the bovine  $\alpha$ S1-casein gene.

In an alternative embodiment a genomic sequence, such as a human genomic clone or clones, can be introduced into an animal to produce a transgenic animal containing  
15 a transgene that has the sequence of the human gene, including all or part of the 5' expression regulation sequences, coding sequences, introns, and 3' untranslated and flanking sequences. In a preferred embodiment, the human lactoferrin genomic sequence is  
20 used in its entirety, but various components can be substituted with components from other mammary gland specific genes.

Although the use of both 5' and 3' expression regulation sequences are preferred, in some embodiments of the  
25 invention, endogenous 3' regulation sequences are not used. In such cases, the 3' proximal expression regulation sequences normally associated with the genomic DNA encoded by the recombinant DNA sequence are used to direct polyadenylation. In addition, distal 3'  
30 regulation sequences from the genomic DNA encoding the recombinant polypeptide may also be employed preferably in the same amounts as set forth for endogenous 3' expression regulation sequences. In such cases, it is to be understood that the recombinant polypeptide

-22-

encoded by the transgene may comprise either genomic DNA or a double stranded DNA derived from cDNA. As with the 5' expression regulation sequences, the optimal amount of 3' expression regulation sequence may be readily  
 5 determined by varying the amount of 3' flanking sequence to obtain maximal expression of the recombinant polypeptide. In general, the distal 3' regulation sequence, be it from an endogenous gene or a heterologous gene, will not extend into the adjacent  
 10 gene from which is derived and will exclude any sequences which adversely effect the level of transgene expression.

Examples of expression regulation sequences are provided in Table I.

15

TABLE 1

	<u>Expression Regulation Sequence</u>	<u>Tissue Specificity</u>	<u>Animal Species</u>
20	16 kb of bovine $\alpha$ S1 casein 5' to structural gene and 8 kb 3' to structural gene	Mammary secretory cells	bovine
	$\approx$ 15 kb 5' to albumin gene	Liver	murine
25	$\approx$ 15 kb 5' to $\alpha$ -actin gene	Muscle	murine
	$\approx$ 15 kb upstream of protamine gene	Spermatids	murine

In addition to the 5' and 3' expression regulation sequences and the recombinant DNA (either genomic or  
 30 derived from cDNA) the transgenes of the invention preferably also comprise a "recombinant intervening sequence" which interrupts the transcribed but untranslated 5' region of the transgene. Such intervening sequences can be derived, for example, from



-23-

bovine  $\alpha$ S1 casein and from human lactoferrin. Such sequences as used herein are "homologous recombinant intervening sequences" in that the 5' and 3' RNA splice signals in such recombinant intervening sequences are those normally found in an intervening sequence from an endogenous or heterologous gene. Recombinant intervening sequences may, however, also comprise a "hybrid intervening sequence". Such hybrid intervening sequences comprise a 5' RNA splice signal and 3' RNA splice signal from intervening sequences from different sources. In some aspects of the invention, such hybrid intervening sequences comprise at least one "permissive RNA splice sequence". As used herein, a permissive RNA splice signal is an RNA splice signal sequence, preferably a 3' RNA splice signal, from an intron contained within a repertoire of germ line DNA segments which undergo rearrangement during cell differentiation. Examples of such gene repertoires include the immunoglobulin super gene family, including the immunoglobulins and T-cell antigen receptors as well as the repertoire of the major histocompatibility complex (MHC) genes and others. Particularly preferred permissive splice sequences are those obtained from the immunoglobulin repertoire, preferably of the IgG class, and more preferably those 3' splice signal sequences associated with the J-C segment rearrangement of the Ig heavy and light chain, most preferably the heavy chain. A particularly preferred permissive splice sequence comprises that portion of the sequence as shown downstream of the HindIII site in Fig. 11. A particularly preferred hybrid intervening sequence comprises the entire sequence shown in Fig. 11 which includes a 5' portion of an intervening sequence from bovine  $\alpha$ S1 casein and a 3' sequence portion of an IgG heavy chain intervening sequence.

-24-

Such hybrid intervening sequences containing permissive RNA splice signals are preferably used when the recombinant DNA corresponds to a cDNA sequence. As indicated in the Examples, when 16 kb of 5' expression regulation sequence from the  $\alpha$ S1 casein gene was used in conjunction with an  $\alpha$ S1 casein-IgG hybrid intervening sequence to express human lactoferrin cDNA operably linked to the  $\alpha$ S1 casein secretory signal sequence a transgenic mouse was obtained which produced approximately 1330  $\mu$ g/ml of hLF in the transgenic milk. This amount of recombinant polypeptide far exceeds the previously reported amounts for production of various protein in transgenic mouse milk of generally less than 10  $\mu$ g/ml and in one case approximately 50  $\mu$ g/ml. It also exceeds the maximum of 8  $\mu$ g/ml of hLF produced herein when the same transgene was used that contained a homologous bovine intervening sequence rather than the hybrid intervening sequence.

However, such hybrid intervening sequences are not limited to transgenes utilizing cDNA sequence. Rather, hybrid intervening sequences are also useful when the recombinant polypeptide is encoded by a genomic sequence. Based on the results obtained with the cDNA recombinant DNA and the general expectation that genomic DNA sequences express at higher levels than sequences derived from cDNA, it is expected that such hybrid intervening sequences used in conjunction with genomic recombinant DNA will further enhance expression levels above that which would otherwise be obtained with genomic sequence alone.

Based on the foregoing, it is apparent that preferred transgenes include large amounts of 5' and 3' expression regulation sequences. Further, the recombinant DNA is preferably derived from genomic clones which may be tens to hundreds of kilobases in length. Based on the

-25-

present technology for cloning and manipulating DNA, the construction and microinjection of transgenes is practically limited to linearized DNA having a length not greater than about 50 kb. However, the transgenes of the invention, especially those having a length greater than about 50 kb, may be readily generated by introducing two or more overlapping fragments of the desired transgene into an embryonal target cell. When so introduced, the overlapping fragments undergo homologous recombination which results in integration of the fully reconstituted transgene in the genome of the target cell. In general, it is preferred that such overlapping transgene fragments have 100% homology in those regions which overlap. However, lower sequence homology may be tolerated provided efficient homologous recombination occurs. If non-homology does exist between the homologous sequence portions, it is preferred that the non-homology not be spread throughout the homologous sequence portion but rather be located in discrete areas. Although as few as 14 base pairs at 100% homology are sufficient for homologous recombination in mammalian cells (Rubnitz, J. and Subramani, S. (1984) Mol. Cell. Biol. 4:2253-2258), longer homologous sequence portions are preferred, e.g. 500bp, more preferably 1000bp, next most preferably 2000bp and most preferably greater than 2000bp for each homologous sequence portion.

As indicated in the examples, three overlapping fragments of the human serum albumin gene were microinjected into the pronuclei of mouse zygotes in approximately equal molar portions. These fragments successfully recombined and integrated into the mouse genome as confirmed by analysis of the integrated DNA by Southern blotting procedures and by detection of RNA transcript and human serum albumin in the serum of the transgenic mouse. Although the transgene so generated

-26-

has a unit length of 38 kb, there is no known practical limit to the size of the transgene which may be formed using larger and/or greater numbers of overlapping transgene fragments. In particular, it is expected that

5 transgenes may be formed by this approach having lengths between about 50 to 1000 kb and more preferably between 50 and 500 kb. Further, the use of homologous recombination of overlapping fragments is expected to be fruitful in the generation of larger transgenic

10 animals, such as transgenic bovine species, containing transgenes incorporating recombinant DNA comprising genomic DNA which otherwise could not be incorporated into a pronucleus to form a transgenic animal. Such genomic transgenes are expected to produce higher

15 expression levels in transgenic cows as compared to that which is produced by transgenes encoding recombinant cDNA.

When, the ultimate object is to secrete a recombinant polypeptide, a "secretory DNA sequence" encoding a

20 functional secretion signal peptide is also operably linked within the transgene to direct secretion of the recombinant polypeptide from one or more cell types within the transgenic animal. Secretory DNA sequences in general are derived from genes encoding secreted

25 proteins of the same species of the transgenic animal. Such secretory DNA sequences are preferably derived from genes encoding polypeptides secreted from the cell type targeted for tissue-specific expression, e.g. secreted milk proteins for expression in and secretion from

30 mammary secretory cells. Secretory DNA sequences, however, are not limited to such sequences. Secretory DNA sequences from proteins secreted from other cell types within the species of transgenic animal may also be used, e.g., the native signal sequence of a

35 homologous gene encoding a protein secreted other than in the mammary glands. In addition, "heterologous

-27-

secretory DNA sequences" which encode signal secretion peptides from species other than the transgenic animals may also be used e.g., human t-PA, human serum albumin human lactoferrin and human lactalbumin and secretion

5 signals from microbial genes encoding secreted polypeptides such as from yeast, filamentous fungi, and bacteria. In general, a secretory DNA sequence may be defined functionally as any DNA sequence which when operably linked to a recombinant DNA sequence encodes

10 a signal peptide which is capable of causing the secretion of the recombinant polypeptide.

In one of the preferred embodiments, a secretory DNA sequence encoding a secretory signal sequence functional in the mammary secretory cells of bovine species is used

15 to cause secretion of recombinant polypeptide from bovine mammary secretory cells. The secretory DNA sequence is operably linked to the recombinant DNA sequence. Examples of such secretory DNA sequences include DNA sequences encoding signal secretion

20 sequences for bovine  $\alpha$ S1 casein, murine lactoferrin and human transferrin. The preferred secretory DNA sequence is that encoding the secretory sequence of  $\alpha$ S1 casein from bovine species. The use of this secretory DNA sequence is described in more detail in the Examples.

25 "Operably linked" in the context of linking a secretory DNA sequence to a recombinant DNA sequence means that the secretory DNA sequence (comprising codons encoding the secretory signal peptide sequence) is covalently coupled to the recombinant DNA sequence so that the

30 resultant secretory-recombinant DNA sequence encodes 5' to 3' for the secretory signal sequence and recombinant polypeptide. Accordingly, the reading frame for the secretory sequence and the recombinant DNA sequence must be covalently combined such that an open reading frame

35 exists from the 5' end of the mRNA sequence formed after

-28-

transcription and processing of the primary RNA transcript. This open reading frame in the RNA contains a 5' sequence portion encoding the secretory signal peptide and a 3' sequence portion encoding the recombinant polypeptide. When so constructed, the recombinant polypeptide produced upon expression of the secretory-recombinant DNA sequence is of a form which is capable of being secreted from targeted cells which express the DNA sequence. The signal peptide generally is removed in vivo during secretion to produce an extracellular form of the recombinant polypeptide.

In the preferred embodiments of the invention, a secretory-recombinant DNA sequence is expressed predominantly in the mammary secretory cells of transgenic bovine species. Such tissue-specific expression is obtained by operably linking mammary specific expression regulation DNA sequences to the above secretory-recombinant DNA sequence. Such mammary specific regulation sequences include the aforementioned regulation sequences contained in various bovine genes preferentially expressed in the mammary secretory cells of the species. Such mammary specific genes include  $\alpha$ S1 casein;  $\alpha$ S2-casein;  $\beta$ -casein; K-casein;  $\alpha$ -lactalbumin; and  $\beta$ -lactoglobulin. Preferred expression regulation sequences are derived from  $\alpha$ S1 casein as described more in detail in the Examples.

In general, the transgenes of the invention that are designed to secrete the recombinant polypeptide into transgenic bovine milk are capable of causing such secretion at levels significantly higher than that previously reported for transgenic mice and sheep. When the recombinant polypeptide is encoded by a recombinant DNA corresponding to, or derived from, cDNA, the molar concentration of the recombinant polypeptide is preferably greater than about 1.0  $\mu$ M, more preferably

-29-

greater than about 100  $\mu\text{M}$ , and most preferably greater than 100  $\mu\text{M}$ . When viewed from the perspective of the level of recombinant polypeptide present in the transgenic milk, the amount of recombinant polypeptide  
5 is preferably greater than 50  $\mu\text{g/ml}$ , more preferably greater than about 500  $\mu\text{g/ml}$  and most preferably greater than about 1000  $\mu\text{g/ml}$  (1mg/ml).

When the transgene of the invention encodes a recombinant polypeptide that is encoded by recombinant  
10 DNA derived from or corresponding to genomic DNA (or comprised substantially of such genomic sequences, e.g. greater than about 50%, more preferably greater than about 75%, most preferably greater than 90% of the codons encoding the recombinant polypeptide are from  
15 genomic sequences), the molar concentrations and protein levels in bovine transgenic milk are the same as for cDNA or higher. In general, the molar concentration of the recombinant polypeptide in such transgenic milk is preferably greater than about 50  $\mu\text{M}$ , more preferably  
20 greater than about 150  $\mu\text{M}$ , most preferably greater than about 500  $\mu\text{M}$ . When viewed from the level of protein in the transgenic milk, the levels are preferably greater than about 10 mg/ml, more preferably greater than about 2.5 mg/ml, most preferably greater than 5 mg/ml.

25 The foregoing molar concentration and protein levels in bovine transgenic milk will vary depending upon the molecular weight of the particular recombinant polypeptide. A particular advantage of producing a recombinant polypeptide in bovine transgenic milk is  
30 that relatively large molecular weight polypeptides may be so produced which are otherwise difficult to produce in large quantities in other systems such as prokaryotic expression systems. Although any recombinant polypeptide may be produced in bovine transgenic milk  
35 according to the invention, it is generally preferred

-30-

that such recombinant polypeptides have a molecular weight greater than about 10,000 Daltons. However, other recombinant polypeptides having molecular weights of greater than 15,000, greater than 20,000 and greater than 60,000 Daltons may also be expressed in transgenic bovine milk. For example, human lysozyme having a molecular weight of 17,000 Daltons and lactoferrin having a molecular weight of 79,000 Daltons may be readily produced in the transgenic milk of bovine species according to the disclosure of the invention. Thus, the recombinant polypeptides of the invention have a wide range of molecular weights.

As a consequence, the foregoing preferred molar concentrations of recombinant polypeptides are adjusted when higher molecular weight recombinant polypeptides are produced. Such adjustment is made by converting the molar concentration to the amount of protein produced and adjusting the molar concentrations so that the recombinant protein level is within the following preferred concentrations.

Most of the previous reports relating to the production of polypeptides in transgenic milk involve transgenic mice. The mouse, however, normally produces between 55 to 80 milligrams of protein per ml of milk. A cow, on the other hand, normally produces between 30 to 34 milligrams of protein per ml. Since exceptionally high levels of recombinant polypeptide production may adversely affect the production of endogenous milk protein and/or have adverse effects upon the mammary secretory gland, it is preferred that the recombinant polypeptide concentration be between about 3 and 50% of the normal bovine milk protein concentration (*i.e.*, between about 1 and 17 milligrams of recombinant polypeptide per ml of transgenic milk), more preferably between 10 to 20% (*i.e.*, between 3 to about 7 milligrams



-31-

per ml) and most preferably between 10 and 15% (i.e., between about 3 and 5 milligrams per ml) of the normal amount of protein produced in bovine milk. Such preferred ranges also provide a preferred maximum limit to the aforementioned levels of protein produced in transgenic bovine milk.

The above described linking of various DNA sequences to form the transgene of the invention are performed by standard methods known to those skilled in the art or as described herein. Once the transgene or overlapping homologous fragments encoding the transgene are constructed as described they are used to make transgenic non-human animals.

Methods of introducing transgenes or overlapping transgene fragments into embryonal target cells include microinjection of the transgene into the pronuclei of fertilized oocytes or nuclei of ES cells of the non-human animal. ~~Such methods for murine species are well known to those skilled in the art.~~ Alternatively, the transgene may be introduced into an animal by infection of zygotes with a retrovirus containing the transgene (Jaenisch, R. (1976) Proc. Natl. Acad. Sci. USA 73:1260-1264). The preferred method is microinjection of the fertilized oocyte. In this preferred embodiment, the fertilized oocytes are first microinjected by standard techniques. They are thereafter cultured in vitro until a "pre-implantation embryo" is obtained. Such pre-implantation embryos preferably contain approximately 16 to 150 cells. The 16 to 32 cell stage of an embryo is commonly referred to as a morula. Those pre-implantation embryos containing more than 32 cells are commonly referred to as blastocysts. They are generally characterized as demonstrating the development of a blastocoel cavity typically at the 64 cell stage. Methods for culturing fertilized oocytes to the pre-

-32-

implantation stage include those described by Gordon, et al. (1984) Methods in Enzymology 101:414; Hogan, et al. (1986) in Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (for the mouse embryo); and Hammer, et al. (1985) Nature 315:680 (for rabbit and porcine embryos) Gandolfi, et al. (1987) J. Reprod. Fert. 81:23-28; Rexroad, et al. (1988) J. Anim. Sci. 66:947-953 (for ovine embryos) and Eyestone, W.H. et al. (1989) J. Reprod. Fert. 85:715-720; Camous., et al. (1984) J. Reprod. Fert. 72:779-785; and Heyman, Y., et al. (1987) Theriogenology 27:5968 (for bovine embryos). Such pre-implantation embryos are thereafter transferred to an appropriate female by standard methods to permit the birth of a transgenic or chimeric animal depending upon the stage of development when the transgene is introduced. As is well known, mosaic animals can be bred to form true germline transgenic animals.

Since the frequency of transgene incorporation is often low, the detection of transgene integration in the pre-implantation embryo is highly desirable. In one aspect of the invention methods are provided for identifying embryos wherein transgenesis has occurred and which permit implantation of transgenic embryos to form transgenic animals. In this method, one or more cells are removed from the pre-implantation embryo. When equal division is used, the embryo is preferably not cultivated past the morula stage (32 cells). Division of the pre-implantation embryo (reviewed by Williams et al. (1986) Theriogenology 22:521-531) results in two "hemi-embryos" (hemi-morula or hemi-blastocyst) one of which is capable of subsequent development after implantation into the appropriate female to develop in utero to term. Although equal division of the pre-implantation embryo is preferred, it is to be understood that such an embryo may be unequally divided either

-33-

- intentionally or unintentionally into two hemi-embryos which are not necessarily of equal cell number. Essentially, all that is required is that one of the embryos which is not analyzed as hereinafter described
- 5 be of sufficient cell number to develop to full term in utero. In a specific embodiment, the hemi-embryo which is not analyzed as described herein, if shown to be transgenic, is used to generate a clonal population of transgenic non-human animals.
- 10 One of each of the hemi-embryos formed by division of pre-implantation embryos is analyzed to determine if the transgene has been integrated into the genome of the organism. Each of the other hemi-embryos is maintained for subsequent implantation into a recipient female of
- 15 the species. A preferred method for detecting transgenesis at this early stage in the embryo's development uses these hemi-embryos in connection with a unique property of the restriction endonuclease Dpn I. This enzyme recognizes the sequence GATC in double-
- 20 stranded DNA but only when the adenine in each strand within this sequence is methylated at N-6. When using this preferred method, the transgene containing the sequence GATC is methylated prior to microinjection either by transferring the transgene on an appropriate
- 25 plasmid through a DAM<sup>+</sup> strain of microorganisms such as E. coli MM294 or by directly methylating the transgene with dam methylase. The methylated transgene (preferably without any exogenous sequences such as plasmid vector) is then microinjected into fertilized
- 30 oocytes (approximately 10 to 500 copies per pronucleus, more preferably 50 to 100 copies per pronucleus). The fertilized oocytes so obtained are cultured in vitro to the pre-implantation stage. During this early growth and cell division phase, the genomic DNA is replicated.
- 35 Accordingly, those copies of the methylated transgene integrated into the genome of the fertilized oocyte are

-34-

unmethylated after replication whereas any non-integrated transgenes which may still exist after replication will remain methylated. (Lacks, S., et al. (1977) J. Mol. Biol. 114:153.) This differential  
5 methylation pattern for integrated versus non-integrated transgene permits the identification of which fertilized oocytes have integrated the transgene into the genome.

The identification of the pre-implantation embryos containing the integrated transgene is achieved by  
10 analyzing the DNA from each of the hemi-embryos. Such DNA is typically obtained by lysing the hemi-embryo and analyzing the thus released DNA after treatment as described by Ninomiya, T. et al. (1989) Molecular Reproduction and Development 1:242-248. Each of the DNA  
15 samples is treated with Dpn I. Thereafter, a polymerase chain reaction (Saiki, et al. (1985) Science 230:1350-1354) is performed to amplify all or part of the transgene. When the entire transgene is amplified, two extension primers each complementary to opposite strands  
20 at opposing ends of the transgene are used for amplification. When, however, less than the entire transgene is amplified, such extension primers are chosen such that the amplified gene product spans the Dpn I site in the transgene. If Dpn I cleavage has not  
25 occurred, PCR amplification results in amplified sequences having a predetermined size whereas primer extension for those transgenes which have been cleaved will not result in exponential amplification. Generally, the Dpn I/PCR amplified DNA from the hemi-  
30 embryo is subjected to electrophoresis followed by hybridization with labeled probe complementary to the region of the transgene between the two extension primers. This facilitates the determination of the size of the amplified DNA sequences, if any, and provides an  
35 indication of whether the transgene has been integrated into the pre-implantation embryo from which the hemi-

-35-

embryo was obtained (now called a "transgenic hemi-embryo"). If it has, the remaining untreated transgenic hemi-embryo is transplanted into a recipient parent. After in utero development, the transgenic non-human  
5 animal having the desired phenotype conferred by the integrated transgene is identified by an appropriate method in utero or after birth. Of course, other restriction endonucleases capable of cleaving a methylated DNA sequence but incapable of cleaving the  
10 unmethylated form of a recognition sequence may be used in the aforementioned method.

The above described method using Dpn I requires that the sequence GATC be present in the transgene of interest. In those cases when such a sequence is not present, it  
15 may be readily introduced into the transgene by site directed mutagenesis (Kunkel, T.A. (1985) Proc. Natl. Acad. Sci. 82:488) or cassette mutagenesis (Wells, J.A., et al. (1985) Gene 34:315) provided such mutagenesis does not change the amino acid sequence encoded by the  
20 transgene (or causes an inconsequential change in amino acid sequence) and that any codons so generated are functional in the transgenic non-human animal of interest.

The above described methods for the detection of  
25 transgenesis in pre-implantation embryos provide economical and time saving method for generating transgenic non-human animals since they significantly decrease the number of pregnancies required to produce a transgenic animal and substantially increase the  
30 likelihood that an implanted embryo will produce a transgenic non-human animal. Such methods are especially important for those animals for which very low or non-existent frequencies of transgenesis have been obtained, e.g. bovine species.

-36-

In an alternate embodiment, the above described method for detecting transgenesis in pre-implantation embryos is combined with embryonic cloning steps to generate a clonal population of transgenic embryos which may thereafter be implanted into recipient females to produce a clonal population of transgenic non-human animals also having the same genotype. In this regard, it is to be understood that transgenic embryos and/or non-human transgenic animals having the same "genotype" means that the genomic DNA is substantially identical between the individuals of the embryo and/or transgenic animal population. It is to be understood, however, that during mitosis various somatic mutations may occur which may produce variations in the genotype of one or more cells and/or animals. Thus, a population having the same genotype may demonstrate individual or subpopulation variations.

After a hemi-embryo is identified as a transgenic hemi-embryo, it is cloned. Such embryo cloning may be performed by several different approaches. In one cloning method, the transgenic hemi-embryo is cultured in the same or in a similar media as used to culture individual oocytes to the pre-implantation stage. The "transgenic embryo" so formed (preferably a transgenic morula) is then divided into "transgenic hemi-embryos" which can then be implanted into a recipient female to form a clonal population of two transgenic non-human animals. Alternatively, the two transgenic hemi-embryos obtained may be again cultivated to the pre-implantation stage, divided, and recultivated to the transgenic embryo stage. This procedure is repeated until the desired number of clonal transgenic embryos having the same genotype are obtained. Such transgenic embryos may then be implanted into recipient females to produce a clonal population of transgenic non-human animals.

-37-

In a preferred cloning method, the transgenic embryo is cloned by nuclear transfer according to the techniques of Prather, et al. (1988) Biol. Reprod. 37:59-86; Roble, et al. (1987) J. Anim. Sci. 64:642-664. According to  
5 this method, nuclei of the transgenic embryo are transplanted into enucleated oocytes, each of which is thereafter cultured to the blastocyst stage. At this point, the transgenic embryos may be resubjected to another round of cloning by nuclear transplantation or  
10 may be transferred to a recipient parent for production of transgenic offspring having the same genotype.

In addition to the foregoing methods for detecting early transgenesis, other methods may be used to detect transgenesis. Such methods include in utero and post  
15 partum analysis of tissue. In utero analysis is performed by several techniques. In one, transvaginal puncture of the amniotic cavity is performed under echoscopic guidance (Bowgso, et al. (1975) Bet. Res. 96:124-127; Rumsey, et al. (1974) J. Anim. Sci.  
20 39:386-391). This involves recovering about 15 to 20 milliliters of amniotic fluid between about day 35 and day 100 of gestation. This volume of amniotic fluid contains about 1000 to 12,000 cells per ml originating from the urogenital tract, the skin and possibly the  
25 lungs of the developing embryo. Most of these cells are dead. Such cells, however, contain genomic DNA which is subjected to PCR analysis for the transgene as an indication of a successful transgenesis. Alternatively, fetal cells may be recovered by chorion puncture. This  
30 method also may be performed transvaginally and under echoscopic guidance. In this method, a needle is used to puncture the recipient animal's placenta, particularly the placentonal structures, which are fixed against the vaginal wall. Such sampling may be  
35 performed around day 60 of gestation in bovine species. Chorion cells, if necessary, are separated from maternal

-38-

tissue and subjected to PCR analysis for the transgene as an indication of successful transgenesis.

Transgenesis may also be detected after birth. In such cases, transgene integration can be detected by taking  
5 an appropriate tissue biopsy such as from the ear or tail of the putative transgenic animal. About one to two centimeters of tail or about five to ten square millimeters of ear are obtained followed by southern blotting with a probe for the transgene according to the  
10 method of Hogan, et al. (1986) Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory.

Transgenesis can also be determined by using the southern blot technique with DNA obtained from other tissues. In particular, semen from a recombinant bull  
15 will be useful for identifying transgenic animals.

Transgenesis may also be detected by assaying for expression of the recombinant polypeptide in a tissue, secretion (e.g., saliva), or other body fluid. In the case where the goal is expression of a recombinant  
20 polypeptide in milk of cows it will be especially useful to assay the saliva of bulls for expression levels. This is because some mammary specific promoters may also cause salivary gland expression, albeit at low levels. See, e.g., Archibald et al. (1990) Proc. Nat. Acad. Sci.  
25 USA 87Z:5178-5182.

In those embodiments where a recombinant polypeptide is expressed and secreted into the milk of transgenic bovine species, the transgenic milk so obtained may be either used as is or further treated to purify the  
30 recombinant polypeptide. This depends, in part, on the recombinant polypeptide contained in the transgenic milk and the ultimate use for that protein. Thus, when the recombinant polypeptide is secreted into transgenic milk



-39-

to increase the nutritional value of the bovine milk, no further purification is generally necessary. An example of such a situation involves one of the preferred embodiments wherein human lactoferrin is produced in the milk of bovine species as a supplement to control intestinal tract infections in newborn human infants and to improve iron absorption. In other situations, a partial purification may be desired to isolate a particular recombinant polypeptide for its nutritional value. Thus, for example, human lactoferrin produced in transgenic bovine milk may be partially purified by acidifying the milk to about pH 4-5 to precipitate caseins. The soluble fraction (the whey) contains the human lactoferrin which is partially purified.

The recombinant polypeptide contained in bovine transgenic milk may also be used in food formulations. A particularly useful food formulation comprises an infant formula containing one or more recombinant polypeptides from transgenic bovine milk which have either nutritional or other beneficial value. For example, an infant formula containing human lactoferrin from transgenic bovine milk made according to the present invention provides a bacteriostatic effect which aids in controlling diarrhea in newborn. Similarly, recombinant polypeptides such as human casein and human lysozyme may also be generated in transgenic bovine milk to provide nutritional value. Table 2 sets forth the constituents of a typical infant formula. As indicated therein, the protein content varies between about 1.8 and 4.5 grams of protein per 100 kilocalories of formula. Thus, the total protein including recombinant polypeptide should lie between the values at least based on regulatory requirements in the United States from which the formulation in Table 2 is based. The amount of total protein including recombinant polypeptide, of

-40-

course, may vary from the foregoing depending upon the local regulations where the particular formula is intended to be used.

-41-

TABLE 2

Nutrient	Minimum <sup>a</sup>	Maximum <sup>a</sup>
Protein (gm)	1.8 <sup>a</sup>	4.5
Fat:		
gm	3.3	6.0
percent cal	30.0	54.0
Essential fatty acids (linoleate):		
percent cal	2.7	
mg	300.0	
Vitamins:		
(A) (IU)	250.0 (75 µg)	750.0 (225 µg) <sup>c</sup>
D (IU)	40.0	100.0
K (µg)	4.0	
E (IU)	0.7 (with 0.7 IU/gm linoleic acid)	
C (ascorbic acid (mg)	8.0	
B <sub>1</sub> (thiamine (µg)	40.0	
B <sub>2</sub> (riboflavin) (µg)	60.0	
B <sub>6</sub> (pyridoxine) (µg)	35.0 (with 15 µg/gm of protein in formula)	
B <sub>12</sub> (µg)	0.15	
Niacin (µg)	250.0	
Folic acid (µg)	4.0	
Pantothenic acid (g)	300.0	
Biotin (µg)	1.5 <sup>a</sup>	
Choline (mg)	7.0 <sup>a</sup>	
Inositol (mg)	4.0 <sup>a</sup>	
Minerals:		
Calcium (mg)	50.0 <sup>a</sup>	
Phosphorus (mg)	25.0 <sup>a</sup>	
Magnesium (mg)	6.0	
Iron (mg)	0.15	
Iodine (µg)	5.0	
Zinc (mg)	0.5	
Copper (µg)	60.0	
Manganese (µg)	5.0	
Sodium (mg)	20.0	60.0
Potassium (mg)	80.0	200.0
Chloride (mg)	55.0	150.0

<sup>a</sup>Stated per 100 kilocalories.<sup>b</sup>The source of protein shall be at least nutritionally equivalent to casein.<sup>c</sup>Retinol equivalents.<sup>d</sup>Required to be included in this amount only in formulas which are not milk-based.<sup>e</sup>Calcium to phosphorus ratio must be no less than 1.1 nor more than 2.0.<sup>f</sup>Includes recombinant protein according to the invention or recombinant proteins and other proteins.

-42-

In addition to infant formulas, other food formulations may also be supplemented with recombinant polypeptides from transgenic bovine milk. For example, such recombinant polypeptides may be used to supplement  
5 common diet formulations.

When the recombinant polypeptide is intended to be used pharmaceutically, purification methods consistent with such an application are called for. Such purification methods will depend on the particular recombinant  
10 polypeptide to be purified and are generally known to those skilled in the art. Such methods typically include a partial purification by casein fractionation followed by chromatography of the appropriate fraction containing the recombinant polypeptide. Such  
15 chromatography includes affinity chromatography, ion exchange chromatography, gel filtration and HPLC.

In a specific embodiment of the invention, transgenes are provided for producing human lactoferrin in the milk of transgenic bovine species. Human lactoferrin (HLF)  
20 is a single chain glycoprotein which binds two ferric ions. Secreted by exocrine glands (Mason, et al. (1978) J. Clin. Path. 31:316-327; Tenovuo, et al. (1986) Infect. Immun. 51:49-53) and polymorphonuclear neutrophil granulocytes (Mason, et al. (1969) J. Exp.  
25 Med. 130:643-658), this protein functions as part of a host non-specific defense system by inhibiting the growth of a diverse spectrum of bacteria. HLF exhibits a bacteriostatic effect by chelation of the available iron in the media, making this essential metal  
30 inaccessible to the invading microorganisms (Bullen, et al. (1972) Br. Med. J. 1:69-75; Griffiths, et al. (1977) Infect. Immun. 15:396-401; Spik, et al. (1978) Immunology 8:663-671; Stuart, et al. (1984) Int. J. Biochem. 16:1043-1947). This effect is blocked if the  
35 protein is saturated with ferric ions. Several studies

-43-

suggest that HLF displays a direct bacteriocidal effect on certain microorganisms (Arnold, et al. (1980) Infect. Immun. 28:893-898; Arnold, et al. (1977) Science 197:263-265; Arnold, et al. (1981) Infect. Immun. 5 32:655-660; Arnold, et al. (1982) Infect. Immun. 35:792-797; Bortner, et al. (1986) Infect. Immun. 51:373-377). The bacteriocidal effect is also inhibited by iron saturation of the protein. No mechanism for the bacteriocidal effect of HLF has been postulated, although 10 it has been demonstrated that it can damage the outer membrane and alter outer membrane permeability in gram-negative bacteria (Ellison, et al. (1988) Infect. Immun. 56:2774-2781).

Lactoferrin is the major iron binding protein in human 15 milk (present at a concentration of about 1.5-1.7 mg/ml) and may play a role in the absorption of iron by the small intestine. All of the iron present in breast milk is thought to be bound to hLF and is taken up at very high efficiencies compared to formula (Hide, D.W., et 20 al. (1981) Arch. Dis. Child. 56:172). It has been postulated that the high uptake of the hLF bound iron is due to a receptor in the jejunum and data has been presented suggesting existence of receptors in Rhesus monkeys (Cox, et al. (1979) BBA 588:120; Davidson, L.A., 25 et al. (1985) Fed. Proc. 18:901). There is also evidence for specific lactoferrin receptors on mucosal cells of the small intestine of human adults (Cox, et al. (1979) Biochem. Biophys. Acta. 588:120-128). Free iron levels have been implicated in the control of the 30 intestinal flora (Mevissen-Verhage, et al. (1985) Eur. J. Clin. Microbiol. 4:14). Breast fed infants, compared with infants fed cow's milk, with and without added iron, were shown to have substantially reduced coliform and, elevated bifidobacteria and clostridia counts in 35 fecal samples. In in vitro studies, human milk has been shown to have a specific inhibitory effect on E. coli

-44-

(Brock, et al. (1983) Infect. and Immunit. 40:453). Human milk has also been shown to have a specific inhibitory effect on E. coli in small intestine due to its high content of iron binding protein, predominantly hLF (Bullen, et al. (1972) British Med. J. i:69).

Thus, the production of human lactoferrin in the milk of transgenic bovine species provides a source of human lactoferrin. Such lactoferrin may be purified from the transgenic milk for formulation purposes.

10 Alternatively, the whole transgenic milk may be used, preferably after pasteurization, in either liquid or dried form. In addition, the beneficial action of human lactoferrin may be potentiated by combining the human lactoferrin or the transgenic milk containing it with

15 human lysozyme. The human lysozyme may be simultaneously produced in the transgenic cow by introducing a second transgene simultaneously with the HLF transgene to produce a transgenic cow capable of producing more than one recombinant polypeptide in the

20 transgenic milk. Alternatively, the transgenes may be sequentially introduced into bovine species. When such is the case, a transgenic bovine species is obtained containing one of the transgenes. Thereafter, embryonic cells, such as eggs, are obtained from the transgenic

25 female and treated so as to incorporate the second transgene encoding the second polypeptide. Preferably, the egg is fertilized, followed by microinjection of the pronucleus of the zygote so obtained. It is to be understood that the foregoing combination of more than

30 two recombinant polypeptides in transgenic bovine milk is not limited to the aforementioned human lactoferrin and lysozyme combination. Thus, the invention contemplates the production of transgenic bovine species and transgenic milk wherein more than one recombinant

35 polypeptide is produced by such a transgenic animal in the transgenic milk.

-45-

The complete amino acid sequence of HLF has been determined (Metz-Boutigue et al. (1984) Eur. J. Biochem. 1451:659-676). HLF comprises two domains, each containing one iron-binding site and one N-linked glycosylation site. These domains show homology between each other, indicative of an ancestral gene duplication and fusion event. In addition, HLF shares extensive homology with other members of the transferrin family (Metz-Boutigue, supra; Pentecost, et al. (1987) J. Biol. Chem. 262:10134-10139). Location of the amino acids involved in the iron-binding sites has been determined by X-ray crystallography (Anderson et al. (1987) Proc. Natl. Acad. Sci. 84:1769-1773). A partial cDNA sequence for neutrophil HLF was published by Rado, et al. (1987) Blood 70:989-993. There was a >98% agreement between the amino acid sequence deduced from the cDNA and that which was determined by direct analysis of lactoferrin from human milk. The structure of the iron-saturated and iron-free form of human lactoferrin have recently been published. (Anderson, et al., (1989) J. Mol. Biol. 209:711-734; Anderson, et al. (1990) Nature:784-787.)

As used herein, "human lactoferrin" comprises a polypeptide having the amino acid sequence substantially as described by Metz-Boutigue, et al. (1984) Eur. J. Biochem. 1451:659-676 and as set forth in Fig. 2. It is noted, however, that an earlier partial sequence of the human lactoferrin sequence disclosed a number of discrepancies between the published sequence and that obtained herein. Specifically, the following discrepancies exist (amino acid numbering is from the sequence in Figure 1 with DNA position in parenthesis):

-46-

	<u>Amino Acid</u>	<u>Position</u>	<u>In Metz-Boutique</u>
	Arg	122 (418)	Absent
	Thr	130 (442)	Ile
	Gln	151 (505)	Arg
5	Ser	184 (604)	Leu
	Tyr	189 (619)	Lys
	Ser	372 (1169)	TrP
	between Ala and Met	391 (1122)	13 amino acids
10	Cys	403 (1225)	Gly
	Gln	512 (1588)	Glu
	Lys	675 (2077)	Arg

Accordingly, human lactoferrin is also defined by the sequence shown in Figure 1 which combines the sequence differences obtained herein with the published sequence. The term human lactoferrin also includes allelic variations of either of these sequences or recombinant human lactoferrin variants wherein one or more amino acids have been modified by the substitution, insertion or deletion of one or more amino acid residues. In some instances human lactoferrin may be produced in milk with all or part of a secretory signal sequence covalently attached thereto.

As used herein, a "human lactoferrin DNA sequence" is a DNA sequence which encodes human lactoferrin as defined above. Such a human lactoferrin DNA sequence may be obtained from a human mammary gland cDNA library or may be derived from the human genome. Example 2 herein describes the cloning and nucleotide sequence of human lactoferrin derived from a human mammary gland cDNA library. The DNA sequence of this human lactoferrin is shown in Fig. 1 and Fig. 2 and is substantially the same as that described by Rado, et al. (1987) Blood 70:989-993. The construction of plasmids containing an expressible transgene encoding hLF is described in the examples. One of these plasmids is cGP1HLF also sometimes referred to as 16,8HLF3) contains



-47-

a transgene designed for tissue-specific expression in bovine mammary secretory cells.

In a second embodiment of the invention, transgenes are provided for producing human serum albumin in the milk of transgenic bovine species. Human serum albumin is a serum protein which contains 584 amino acid residues (Minghetti, et al. (1986) J. Biol. Chem. 261:6747). It is the most abundant protein in human serum and performs two very important physiological functions. Serum albumin is responsible for about 80% of the total osmolarity of blood and it transports fatty acids between adipose tissues.

Human serum albumin is used primarily to expand plasma volume by restoring osmotic pressure in the circulatory system. Currently, a heat treated serum derived hSA fraction is infused in most shock and trauma victims, including most of the patients undergoing extensive surgery. HSA is presently derived from human blood plasma as a by-product from blood fractionation processes to obtain rare blood proteins such as factor VIII and IX. The recently developed technology of producing such factors by biotechnological means, however, threatens the source of human serum albumin.

As used herein "human serum albumin" comprises a polypeptide having the amino acid sequence substantially as that described by Minghetti, et al., ibid; Lawn, et al. (1981) Nucl. Acids Res. 9:6103. Also included are variations thereof including recombinant human serum albumin variants wherein one or more amino acids have been modified by the substitution, insertion or deletion of one or more amino acid residues (Minghetti, et al. (1986) J. Biol. Chem. 261:6747-6757). In some instances, human serum albumin may be produced in milk by expressing a transgene which contains DNA encoding

-48-

the secretory signal sequence of hSA. Alternatively, human serum albumin may be produced in and secreted from liver cells of a transgenic animal utilizing a completely heterologous transgene comprising human  
5 genomic DNA encoding 5' expression regulation sequences, the human serum albumin secretion signal and structural gene and 3' expression regulation sequences. As indicated in the Examples, transgenes containing this heterologous sequence were formed by in vivo homologous  
10 recombination of overlapping transgene fragments to reconstitute the hSA gene in the transgenic animal. The so formed transgenic animal produced human serum albumin in its circulatory system.

As used herein, a "human serum albumin DNA sequence" is  
15 a DNA sequence which encodes human serum albumin as defined above. Such a human serum albumin DNA sequence may be obtained from  $\lambda$ HAL-HAI,  $\lambda$ HAL-3W and  $\lambda$ HAL-HI4 as described by Urano, et al. (1986) J. Biol. Chem. 261:3244-3251 and Urano, et al. (1984) Gene 32:255-261  
20 and in the Examples herein:

The human serum albumin DNA sequence was cloned as described in Example 10 herein and subsequently manipulated to substitute for the human lactoferrin gene encoded in plasmid cGP1HLF (also referred to as  
25 p16,8HLF4). From this plasmid a transgene is obtained containing 16 kb of the 5' expression regulation sequence of the bovine  $\alpha$ S1 casein gene, human serum albumin DNA sequence and approximately 8 kb of the 3'-flanking region of the  $\alpha$ S1 casein bovine gene. This  
30 transgene is used to microinject fertilized oocytes from bovine species. After early detection of transgenesis, blastocysts containing the hSA transgene are implanted into a recipient female bovine species and brought to term.

-49-

The following is presented by way of example and is not to be construed as any limitation on the scope of the invention.

#### EXAMPLE 1

##### 5                    Construction of a Probe Specific                     for Bovine $\alpha$ S1 Casein Sequences

###### A.            Isolation of Chromosomal DNA

Placental tissue was obtained from the slaughterhouse.  
10    Surrounding connective tissue was removed and pieces of  
      about 30 grams were quickly frozen in liquid N<sub>2</sub>.  
      Chromosomal DNA was isolated as follows: 30 grams of  
      tissue was homogenized (on ice) with 35 ml of Buffer 1  
      containing 300 mM Sucrose; 60 mM KCl; 15 mM NaCl; 60 mM  
15    Tris.HCl pH 8.2; 0.5 mM spermidine; 0.15 mM spermine;  
      2 mM EDTA; 0.5 mM EGTA. 65 ml of icecold buffer 1  
      containing 1% NP40 was added and the mixture was  
      incubated for five minutes on ice. After centrifugation  
      for five minutes at 3000 xg the pellet was rinsed with  
20    buffer 1 containing 1% NP40. After repeating the  
      centrifugation step the pellet was resuspended in 5 ml  
      of buffer 1. 5 ml 0.5 M EDTA was quickly added. Final  
      volume was now 15 ml. 0.15 ml of a 10% SDS solution was  
      added. After mixing, RNase A and T1 were added to final  
25    concentrations of 0.4 mg/ml and 6 u/ml respectively.  
      After incubation at 37°C for three hours, Proteinase K  
      was added to a final concentration of 0.1 mg/ml. This  
      mixture was incubated for 15 hours at 37°C. The mixture  
      was then carefully extracted with phenol. The aqueous  
30    phase was isolated and 1/30 volume of 3M NaOAc pH 5.2  
      and one volume of isopropylalcohol was added. The  
      precipitate (DNA) was rinsed with 70% ethanol and slowly  
      dissolved in 0.5 ml of 10 mM Tris.HCl pH 8.0; 1 mM EDTA,  
      at 4°C.

-50-

B. Amplification of Sequences from the  
5'-flanking Region of the  $\alpha$ S1-casein Gene

Two DNA-primers were synthesized based on the sequence published by Yu-Lee et al., (1986) Nucl. Acids Res. **14**, 1883-1902. Primer 1 was located at position-681 relative to the major transcription initiation site and had the following sequence:

5'-TCC ATG GGG GTC ACA AAG AAC TGG AC-3'.  
(Seq. ID No.: 5)

Primer #2 was located at position +164 relative to the major transcription initiation site and had the following sequence: 5'-TGA AGC TTG CTA ACA GTA TAT CAT AGG-3' (Seq. ID. No.: 6). The first eight nucleotides of this primer are not encoded by the bovine genome, but contain a HindIII restriction site to facilitate subsequent cloning steps. These primers were annealed to the chromosomal DNA and extended in the presence of deoxynucleotides by TAQ-polymerase. After three minutes the mixture was denatured for one minute at 92°C, reannealed at 50°C for 1.5 minutes and again incubated at extension temperature (68°C) for 2 minutes. This cycle was repeated 30 times. After the last cycle DNA was checked for the presence of the expected EcoRI sites. Both the size of the fragment and the presence of EcoRI sites was as expected. The fragment was then treated with Klenow enzyme to repair any overhanging ends, treated with kinase to attach phosphate groups at the ends of the fragment, incubated at 65°C for 10 minutes to inactivate the kinase and klenow enzymes and finally digested with HindIII. This fragment was then subcloned in pUC19 (Yanisch-Perron, et al. (1985), Gene, **33**, 103-109) digested with SmaI and HindIII. Formal proof of the identity of this fragment was obtained by sequencing parts of this subclone (after re-cloning into M13 vector). The determined sequence was identical to

-51-

the published sequence. This probe was then used to screen a bovine genomic library to obtain clones specific for the 5'-flanking region of the  $\alpha$ S1-casein gene.

5 C. Amplification of Sequences from the  
3'-flanking Region of the  $\alpha$ S1-casein Gene

A similar approach was taken as described above. Two primers were designed based on the sequence published by Stewart et al (1984) Nucl. Acids Res. 12, 3895-3907.

- 10 The 5'-primer was located just downstream of the coding sequence starting at position 713 of the cDNA sequence. It had the following sequence:

5'-GAG GGA CTC CAC AGT TAT GG-3' (Seq. ID No.: 7).

- The other primer was located at position 1070 of the  
15 cDNA sequence and had the following sequence: 5'-GCA  
CAC AAT TAT TTG ATA TG-3' (Seq. ID No.: 8). These  
primers were annealed to the chromosomal DNA and the  
region between these primers was amplified as described  
above. The resulting fragment was  $\approx$ 900 bp longer than  
20 expected. Sequence analysis showed that an intervening  
sequence of this size was present between nucleotide 737  
and 738 of the cDNA. The amplified fragment was treated  
with Klenow-polymerase to repair any overhanging ends  
and treated with kinase to attach phosphate groups to  
25 the ends of the fragment. The fragment was then ligated  
into pUC19 previously cut with SmaI.

-52-

D. Screening of a Bovine Phage Library  
for  $\alpha$ S1-casein Flanking Sequences

A bovine genomic library, constructed in EMBL3, was obtained from Dr. M. Groenen, Agricultural University Wageningen, Netherlands, and was screened in the following way. The bacteriophage particle titre was determined on Escherichia coli MB406 a permissive host strain (Stratagene Inc.). For this, several dilutions of the phage stock were made in SM buffer (50 mM Tris.HCl pH 7.5, 100 mM NaCl, 10 mM MgSO<sub>4</sub>, 0.01% gelatin) and mixed with 200  $\mu$ l MB406 (O.D.<sub>550</sub> = 0.9); after 20 minutes at 37°C, 3 ml top agarose (Luria-Bertani medium, 0.8% agarose, 10 mM MgCl<sub>2</sub>) was added and this was plated on LB plates and incubated overnight at 37°C.

Approximately 600,000 phages were then plated by adding the required amount of phage stock to 400  $\mu$ l MB406. The subsequent plating was as described as above. The next step was transfer of the phage to nitrocellulose filters. Plates were placed at 4°C for one hour. Nitrocellulose filters (S&S) were placed on the top agarose layer and exact position was marked. After lifting, the filters were soaked for (1) 30 minutes in denaturation buffer (1.5M NaCl, 0.5M NaOH); (2) 5 minutes in neutralizing buffer (1.5M NaCl, 0.5M Tris.HCl pH 8.0). After rinsing with 2xSSPE (360 mM NaCl, 20 mM NaH<sub>2</sub>PO<sub>4</sub>, 2 mM EDTA), the filters were baked under vacuum at 80°C for two hours.

Prehybridization of the filters was performed in a buffer containing 50% formamide, 5x Denhardt's solution (0.1% Ficoll, 0.1% polyvinylpyrrolidone, 0.1% bovine serum albumin), 5xSSPE, 0.1% SDS and 100  $\mu$ g/ml denatured salmon sperm DNA at 42°C for two hours. Hybridization was performed in same buffer at 42°C overnight in a shaking waterbath. The probe, generated as previously

-53-

described, was labelled using the Random Primed labelling kit from Boehringer Mannheim. After overnight hybridization the filters were washed three times with 2xSSC, 0.1% SDS at room temperature.

- 5 Overnight exposure of Kodak XAR films was performed with amplifying screens (Dupont) at -70°C. Putative positives were plugged out of the plates and put overnight in SM buffer at 4°C. These were plated out as described above and DNA was isolated following the
- 10 plate lysate method (Maniatis, T., et al. (1982), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, N.Y.). 5 ml SM buffer was added to the top agarose layer; after two hours gentle shaking buffer was removed and spun at 4000 rpm at 4°C for 10 minutes.
- 15 Supernatant was transferred to sterile tubes and RNase A and DNaseI (both final concentration 1µg/ml) was added, this was incubated at 37°C for 30 minutes. One volume of a 20% polyethyleneglycol, 2.5 M NaCl solution was added and put on ice for one hour. Centrifugation
- 20 at 4000 rpm for 30 minutes at 4°C left precipitated bacteriophage particles. These were resuspended in 500 µl SM buffer, SDS (final concentration 0.1%) and EDTA (final concentration 5 mM) was added, this was incubated at 68°C for 15 minutes. Protein was removed
- 25 with one phenol and one chloroform extraction step. Precipitation of phage DNA was performed with one volume isopropanol. Phage DNA was washed once with 70% ethanol and dissolved in 50 µl Tris.HCl pH 7.5, 1 mM EDTA buffer.
- 30 Restriction enzyme analysis, agarose gel electrophoresis, transfer of DNA from gel to nitrocellulose filter and Southern blotting were all done according to standard procedures (Maniatis (1982), Molecular Cloning: A Laboratory Manual). Hybridization
- 35 with probes (described hereinafter) was performed

-54-

according to the same procedure as the screening conditions described above.

E. Isolation of Clones Containing  
5'-flanking Region of Bovine S1-casein

5 Three putative clones were identified using the probe and procedures as described above. After another round of screening, clean recombinant bacteriophage was analyzed. Digestion of cloned DNA with SalI, EcoRI and SalI/EcoRI (double digestion) and hybridization with the  
10 probe described above showed identical inserts in all three clones. The insert consisted of an 18 kb (partial Sau3A fragment excised with SalI). Transcriptional orientation in the clone was determined with hybridization of above described restriction fragments  
15 with (1) probe 1 described above, and (2) the NcoI-NsiI fragment of probe 1. This showed a region of about 16 kb upstream of transcription start. Downstream from the transcription start was another 1.9 kbp. Sequencing of part of the latter region showed the presence of exon  
20 2 and part of intron 2 of the bovine  $\alpha$ S1-casein gene. Additional sequencing of the region -103 - +300 confirmed the identity of the clone. The ethidium-bromide pattern of the described restriction fragments also showed the orientation of the clone in the EMBL vector. Subsequent  
25 analysis of the clone with the following restriction enzymes (NcoI, PstI, KpnI, BamHI, HindIII, BglII) resulted in the restriction map of 5' flanking region of bovine S1-casein gene as shown in Fig. 3.

F. Isolation of Clones Containing  
30 3'-flanking Region of Bovine  $\alpha$ S1-casein

Duplicate nitrocellulose filters from the initial phage plating used for isolating 5' clones were screened with the 3'  $\alpha$ S1-casein probe using the same hybridization conditions previously described. Eight positive clones  
35 were identified after two rounds of screening. Phage



-55-

DNA was prepared as described. Subsequent restriction digests with SaII, EcoRI, and SaI/EcoRI and Southern hybridization with the 3'  $\alpha$ S1 probe showed identical inserts in seven of the eight clones. One clone  
5 containing an 18.5 kb EcoRI insert was further analyzed with the restriction enzymes BsteII and BamHI. A restriction map of that clone is shown in Fig. 4.

## EXAMPLE 2

### Cloning of Human Lactoferrin Gene

#### 10 A. Materials

Restriction endonucleases, T4 ligase, and T7 polynucleotide kinase were obtained from Boehringer-Mannheim, New England Biolabs, or Bethesda Research Laboratories. Radio-isotopes were purchased from  
15 Amersham. A human mammary gland cDNA library in bacteriophage  $\lambda$ gt11 was obtained from Clontech, Inc., Palo Alto, Calif.

#### B. Isolation of the Human Lactoferrin Gene

The human mammary gland library was screened by standard  
20 plaque hybridization technique (Maniatis, et al. (1982) Molecular Cloning: A Laboratory Manual) with three synthetic oligomers. Two of the oligomers were 30-mers corresponding to the cDNA sequence of Rado et al., supra, at amino acid positions 436-445 and 682-691. The  
25 third was a 21-mer "best guess" probe based on human codon bias and coding for amino acid sequence of HLF between amino acid residues 18 and 24. Respectively, they were:

- (1) 5'-CTTGCTGTGGCGGTGGTTAGGAGATCAGAC-3' (Seq. ID  
30 No.: 9)  
(2) 5'-CTCCTGGAAGCCTGTGAATTCCTCAGGAAG-3' (Seq. ID No.: 10), and  
(3) 5'-ACCAAGTGCTTCCAGTGGCAG-3' (Seq. ID No.: 11).

-56-

The probes were radiolabeled (Crouse et al. (1983) Methods Enzymol. 101, 78-98) and used to screen duplicate filters. Filters were washed at a final stringency of 2 X SSC, 37°C.

5 C. Nucleotide Sequence Analysis

DNA fragments were isolated by use of low-melting agarose (Crouse et al, supra) and subcloned into bacteriophage M13mp18 or M13mp19 (Messing et al. (1983) Methods Enzymol. 101, 20-78). The sequence was  
10 determined using the Sequenase enzyme (modified T7 DNA polymerase) (Tabor et al. (1987) Proc. Natl. Acad. Sci. USA 84, 4767-4771). All reactions were carried out according to the manufacturer's specifications (US Biochemicals). The sequence is shown in Fig. 1. The  
15 hLF sequence was digested with HindIII and EcoRI (present in the surrounding phage sequences) and subcloned into the HindIII and EcoRI site of pUC19 to form pUS119 Lacto 4.1. This clone contained the entire coding sequence of the mature form of hLF, but lacked  
20 the complete signal sequence.

EXAMPLE 3

Construction of bovine  $\alpha$ S1-casein CAT vectors

In order to determine whether the  $\alpha$ S1 casein fragments obtained in Example 1 had promoter and other properties  
25 needed to express a heterologous gene, expression plasmids were constructed containing variable amounts of 5-' and 3'-flanking regions from the  $\alpha$ S1-casein gene. The chloramphenicol Acetyl transferase gene (CAT) was used as a heterologous gene in these vector constructs.  
30 The CAT gene is useful to detect the expression level for a heterologous gene construct since it is not normally present in mammalian cells and confers a readily detectable enzymatic activity (see Gorman, C.N.,

-57-

et al. (1983), Mol. Cell. Biol., 2, 1044-1051) which can be quantified in the cells or animals containing an expressible gene.

A. DNA sequences

5 681 bp of a  $\alpha$ S1-casein promoter plus the first non-coding exon plus approximately 150 bp of the first intervening sequence (IVS) were isolated from a 5'-flanking genomic clone from Example 1 by PCR amplification as an NcoI-HindIII fragment (approximately  
10 830 bp). This fragment is identified as fragment 1 in Fig. 5A. The primer sequences consisted of:

5'-TCCATGGGGGTCACAAAGAACTGGAC-3'

(Seq. ID No.: 12) and

5'-TGAAGCTTGCTAACAGTATATCATAGG-3'

15 (Seq. ID No.: 13)

that were designed from a sequence published by Yu-Lee et al. (1986) Nuc. Acids Res. 14, 1883-1902.

Approximately 1.6 kb (fragment 2, Fig. 5A) of  $\alpha$ S1-casein 3'-flanking sequence was isolated by PCR amplification  
20 from a bovine 3'-flanking genomic clone from Example 1. This region contained the previously described splice within the 3' untranslated region of  $\alpha$ S1-casein gene. Fragment 2 was subcloned into the SmaI site of pUC19. The primer sequences consisted of:

25 5'-GAGGGACTCCACAGTTATGG-3'

(Seq. ID No.: 14) and

5'-GCACACAATTATTTGATATG-3'

(Seq. ID No.: 15)

that were designed from a sequence published by Stewart  
30 et al. (1984) Nucl. Acids Res. 12, 3895-3907.

-58-

A hybrid splicing signal comprising the 3' splice site of an immunoglobulin gene (Bothwell et al. (1981), Cell, 24, 625-637) was synthetically prepared and inserted into pUC18 along with unique restriction sites flanking either side to produce pMH-1. This plasmid is shown in Fig. 6. NcoI and HindIII sites were designed such that ligation with fragment 1 from the bovine 5' genomic clone would result in the functional hybrid splice sequence. See Fig. 11.

- 10 A polyadenylation sequence was obtained from SV40 virus as a BamHI-DraI fragment (fragment 3 in Fig. 5A) isolated from pRSVcat (Gorman, C.M., et al. (1982), Proc. Natl. Acad. Sci., 79, 6777-6781).

A bacterial CAT coding sequence was subcloned into pUC19 as a PstI-BamHI fragment.

#### B. Construction of pS13'5'CAT

Fragment 1 of  $\alpha$ S1-casein promoter was subcloned into pMH-1 (Fig. 6) between the NcoI and HindIII sites to form pMHS15'flank.

- 20 The SV40 polyadenylation sequence (fragment 3) was subcloned as a BamHI-DraI fragment into pUC19 immediately 3' to the 3'  $\alpha$ S1-casein flanking sequence (fragment 2) to form pUC19 3' UTR/SV40. This allowed for the removal of a continuous EcoRI-SalI fragment (containing the 3'-flanking sequence and poly (A) sequence) that was subcloned into pMH-1 to derive pMHS13'UTR (Fig. 5B) which was used later to construct pMHSI 3' UTR hlf which contains sequences encoding human lactoferrin.

- 30 The EcoRI-SalI sequence (fragments 2 and 3) were subcloned into the EcoRI-SalI sites of pMHS15'flank to form pS13'5'flank.

-59-

The PstI-BamHI CAT fragment (fragment 4 in Fig. 5B), after blunting the BamHI site with Klenow, was subcloned into pS13'5'flank (Fig. 5B) between the PstI and SmaI sites to form pS13'5'CAT.

5 C. Construction of pS15'CAT

The CAT fragment (fragment 4 in Fig. 5B, PstI-BamHI) and SV40 polyadenylation fragment (fragment 3 in Fig. 5A, BamHI-DraI) were subcloned into the PstI and SmaI sites of pMHS15'flank to form pS15'CAT (Fig. 5C).

10 D. Assay for CAT Production

Each of these CAT plasmids were transfected into human 293S cells (Graham, F.L., et al. (1977), J. Gen. Virol., 36, 59-72) by the calcium phosphate co-precipitation method (Gorman, C.M., et al. (1983), Science, 221, 551; 15 Graham, F.L., et al. (1973), Virology, 52, 456-467). Cells were harvested 44 hours after transfection and cell extracts were assayed for CAT activity (Gorman, C.M., et al. (1982), Mol. Cell. Biol., 2, 1011; 20 deCrombrughe, B., et al. (1973), Nature [London], 241, 237-251, as modified by Nordeen, S.K., et al. (1987), DNA, 6, 173-178). A control plasmid expressing CAT driven by the Cytomegalovirus Immediate early promoter (Boshart, M., et al. (1985), Cell, 41, 521) was transfected into human 293 S cells to assay for 25 transfected efficiency.

pS13'5'CAT was expressed in these cells at a level which was approximately 30-100 fold lower than the control plasmid, but significantly higher than background. Primer extension analysis indicated that transcription 30 had initiated predominantly in the expected region.

When pS15'CAT was transfected into 293S cells, expression was also detected.

-60-

EXAMPLE 4Bovine  $\alpha$ S1-casein/human lactoferrin  
expression cosmid cGP1HLFA. Construction of DNA Sequences.

- 5 16 kb of bovine  $\alpha$ S1-casein 5'-flanking sequence from Example 1 was isolated from the bovine genomic library (phage GP1) as a SalI-BglII fragment. The BglII site lies at the junction of the first intron and second exon of the  $\alpha$ S1-casein gene.
- 10 Bovine  $\alpha$ S1-casein signal sequence (Stewart et al. (1984) Nucl. Acids Res. 12, 3895) was prepared from synthetic DNA synthesized on a Cylone Plus<sup>®</sup> DNA Synthesizer (Millgen/Biosearch I) and contained the entire signal sequence plus XhoI and Cla I sites attached to the
- 15 5'-end, and NaeI to the 3'-end (fragment 8, Fig. 7B).

- Cleavage of pUC119 Lacto 4.1 with EaeI precisely opened the plasmid at the codon for the first amino acid of mature hLF. Treatment with Klenow was used to fill in the overhanging 5'-end. Further digestion with AccI and
- 20 EcoRI gave two fragments: (a) an EaeI-AccI fragment containing the first 243 bp of mature hLF (fragment 5, Fig. 7C), and (b) a contiguous AccI-EcoRI fragment (fragment 6, Fig. 7C) of 1815 bp that contained all but five terminal codons of the remaining coding sequence.

- 25 A synthetic linker was prepared that contained the last five codons of hLF beginning at the EcoRI site and extending for four bases beyond the stop codon. A KpnI site was added to the 3'-end (fragment 7 in Fig. 7C).

- An 8.5 kb EcoRI 3'-fragment was isolated from the bovine
- 30 genomic library (Fig. 4) containing sequences beginning just downstream of the coding region of  $\alpha$ S1-casein and a BstEII site approximately 350 bp from the 5'-end.

-61-

This fragment was subcloned into pMH-1 at the EcoRI site to form pMH3'E10 (Fig. 7A). A SalI site is adjacent to the 3'-EcoRI site in pMH3'E10.

B. Construction of cGP1HLF

- 5 The hLF 3'-linker (fragment 7, Fig. 7C) was subcloned into the EcoRI-KpnI sites of pMH3'UTR (Fig. 7A) to produce pMH3'UTRhLF2linker (Fig. 7A).

The synthetic bovine  $\alpha$ S1-casein signal sequence (fragment 8) was then subcloned into the XhoI and SmaI  
10 sites of pMH3'UTRhLF2linker to make pS13'hLF1/2L (Fig. 7B).

The two hLF coding fragments (fragments 5 and 6 in Fig. 7C) were subcloned into the NaeI and EcoRI sites of pS13'hLF1/2L (Fig. 7B) to make pS13'UTRhLF (Fig. 7C).

- 15 The large  $\alpha$ S1-casein 3'UTR fragment from pMH3'E10 (Fig. 7A) was isolated as a BstEII-SalI fragment and subcloned into the same sites of pS13'UTRhLF to form pHLF3'10 kb (Fig. 7D).

Cosmid cGP1HLF was prepared from a 3-way ligation (Fig. 7F):  
20

- (1) the 16 kb 5'-flanking sequence from phage GP1 (Example 1, Fig. 3) was modified by attaching two linker adapters. The SalI site at the 5'-end was ligated to a NotI-SalI linker. The BglII site at the  
25 3'-end was ligated to a BglII-XhoI linker;

- (2) the hLF coding region, flanked on the 5'-end by the  $\alpha$ S1-casein signal sequence and on the 3'-end by approximately 8.5 kb of  $\alpha$ S1-casein 3'-flanking sequence, was isolated as a XhoI-SalI fragment from  
30 pHLF3'10 kb. The SalI site at the 5'-end was ligated to a SalI-NotI linker;

-62-

(c) Cosmid pWE15 (Stratagene, Inc.) was linearized with NotI.

Fragments from (a), (b), and (c) were ligated together and transfected into bacteria using commercial lambda packaging extracts (Stratagene, Inc.) to produce cGP1HLF.

#### EXAMPLE 5

##### Bovine $\alpha$ S1-casein/hLF expression plasmids.

###### A. Construction of pS13'5'hLF

10 The HindIII-SalI fragment of pS13'UTRhLF was subcloned into the same sites in pMHS15'flank to form pS13'5'hLF (Fig. 7E). This plasmid contains 681 bp of bovine  $\alpha$ S1-casein promoter sequence, the  $\alpha$ S1-casein/IgG hybrid intron, the  $\alpha$ S1-casein signal sequence, the hLF coding  
15 region, approximately 1.6 kb of  $\alpha$ S1-casein 3'-flanking sequence, and the SV40 late region polyadenylation sequence.

###### B. pS15'hLF

Plasmid pS13'5'hLF (Fig. 7E) was cut with KpnI and BamHI  
20 which border the  $\alpha$ S1-casein 1.6 kb 3'-flanking sequence. The larger vector fragment was purified, made blunt ended with Klenow, and self-ligated to form pS15'hLF.

###### C. Radioimmunoassay for hLF

An immunoglobulin-enriched fraction of ascites fluid of  
25 a monoclonal antibody against human lactoferrin, which does not cross-react with the bovine or murine protein, was prepared by 50% ammonium sulfate precipitation and coupled to CNBr-activated Sepharose 4B (20 mg of protein to 1 g of Sepharose). The Sepharose beads were  
30 suspended (2 mg/ml) in phosphate-buffered saline (PBS; 10 mM sodium phosphate, 0.14 M NaCl containing 10 mM



-63-

EDTA, 0.1% (w/v) Polylorene and 0.02% (w/v)  $\text{NaN}_3$ , pH 7.4. Sepharose suspensions (0.3 ml) were incubated for five hours at room temperature by head-over-head rotation with samples (usually 50 $\mu$ l) in 2-ml polystyrene tubes.

5 Sepharose beads were then washed with saline (five times with 1.5 ml) and incubated for 16 hours at room temperature with 50 $\mu$ l (1kBq) of  $^{125}\text{I}$ -labeled-affinity-purified polyclonal rabbit anti human lactoferrin antibodies, together with 0.5 ml of PBS, 0.1% (w/v)

10 Tween-20. Thereafter the Sepharose was washed again with saline (four times with 1.5 ml) and bound radio activity was measured. Results were expressed as percent binding of the labelled antibodies added. Levels of lactoferrin in test samples were expressed in

15 nanomolar, using purified human milk lactoferrin as a standard (serial dilutions in PBS, 10 mM EDTA, 0.1% (w/v) Tween-20.

Repeated testing of standard on separate occasions revealed that this RiA was highly reproducible, intra-

20 and inter assay coefficients of variation ranged from 5-10%. As little as 0.1 nanogram human lactoferrin is easily detected by this RIA.

#### D. Expression in 293S cells

293S cells were transfected with the above hLF plasmids

25 as described (1 $\mu$ g of a CMV-CAT plasmid was co-transfected as control for transfection efficiency). Forty-four hours after transfection medium was removed from the cells and assayed for hLF as described supra, RNA was isolated as described by Stryker, et al. (1989)

30 EMBO J. 8, 2669. The results can be summarized as follows:

1. Transfection efficiencies are identical for the two hLF plasmids;

-64-

2. hLF is expressed in the cells and secreted into the medium. In both cases, the levels are about 0.4 $\mu$ g/ml medium using about  $3 \times 10^6$  cells

3. The proteins behave identical to hLF in a human milk sample in a dose response assay measuring the amount of  $^{125}\text{I}$ - anti-lactoferrin bound as a function of the amount of sample used.

4. The protein has about the same size (~80kD) as in a human milk sample as judged by Western blotting.

10 5. The hLF RNA produced in the cells has the correct size and its level is similar for both plasmids as judged by Northern - blotting.

These data indicate that these two expression plasmids are able to express hLF. By all standards used so far, 15 the protein is identical to hLF present in human milk. The heterologous signal sequence is functional in that it promotes secretion of the protein from the cells into the medium. Further, the casein regulatory sequences used in these plasmids are able to promote expression 20 of a heterologous gene.

#### EXAMPLE 6

##### In vitro Maturation, Fertilization and Culture of Bovine Oocytes

Immature oocytes are obtained in large quantity 25 (400-600/day) by aspirating follicles of ovaries obtained at abbatoirs. Immature oocytes are cultured for a period in vitro before they are competent to be fertilized. Once "matured", oocytes are fertilized with sperm which has also been matured, or "capacitated" in 30 vitro. The pronuclei of the fertilized oocyte is then injected with the transgene encoding for the expression and secretion of human lactoferrin. Zygotes resulting from this in vitro fertilization and microinjection are

-65-

then cultured to the late morula or blastocyst stage (5-6 days) in medium prepared, or "conditioned" by oviductal tissue. Blastocysts are then transferred non-surgically to recipient cattle for the balance of gestation or analyzed for integration of the transgene as described herein.

In vitro maturation (IVM). Ovaries are obtained immediately after slaughter at local abattoirs and oocytes are recovered. Alternatively, oocytes are obtained from living cattle by surgical, endoscopic, or transvaginal ultrasonic approaches. In all cases, oocytes are aspirated from ovarian follicles (2-10 mm diameter). After washing, oocytes are placed in a maturation medium consisting of M199 supplemented with 10% fetal calf serum, and incubated for 24 hours at 39°C. Sirard et al. (1988) Biol. Reprod. 39, 546-552.

In vitro fertilization (IVF). Matured oocytes are fertilized with either fresh or thawed sperm. Sperm are prepared for fertilization by first obtaining a population of sperm enriched for motility by a "swim-up" separation technique (Parrish et al. (1986) Theriogenology 25, 591-600). Motil sperm are then added to a fertilization media, consisting of a modified Tyrode's solution (Parrish et al. (1986) supra.) supplemented with heparin to induce sperm capacitation (Parrish et al. (1988) Biol. Reprod. 38, 1171-1180). Capacitation constitutes the final sperm maturation process which is essential for fertilization. Sperm and oocytes are co-cultured for 18 hours. A useful feature of this IVF method is that (in the case of frozen sperm) consistent, repeatable results are obtained once optimal fertilization conditions for a particular ejaculate have been defined (Parrish et al. (1986) supra.).

-66-

- In vitro culture (IVC). Conventional culture systems, which support development of murine, rabbit, or human ova, do not support development of bovine embryos past the 8-16 cell stage. This problem has been overcome by pre-conditioning culture media with oviductal tissue. Oviduct-conditioned medium will support bovine embryos past the 8-16 cell stage to the blastocyst stage in vitro (Eyestone and First (1989) J. Reprod. Fert. 85, 715-720).
- 10 Bovine embryos have proved refractory to in vitro culture. This in part stems from the existence of a "block" to cleavage in vitro at the 8-16 cell stage. This block may be alleviated by culturing embryos in the oviducts of rabbits (reviewed by Boland (1984)
- 15 Theriogenology 21, 126-137) or sheep (Willadeen (1982) in: Mammalian Egg Transfer, (E. Adams, ed., pp. 185-210)); Eyestone et al. (1987) Theriogenology 28, 1-7). However, these in vivo alternatives have been less than ideal, in that: (1) they require the
- 20 maintenance of large numbers of recipient animals, (2) they require surgery to gain access to the oviducts for transfer, and a second surgery (or sacrifice) to recover the embryos, (3) all transferred embryos are seldom recovered, and (4) access to embryos during
- 25 culture for observation or treatment is entirely precluded. The lack of in vitro culture systems has hampered the development of various manipulation techniques (such as gene transfer by pronuclear injection) by preventing accumulation of basic
- 30 information of the chronology and ontogeny of bovine development, and by complicating the process of culturing embryos to a stage compatible with non-surgical embryo transfer and cryopreservation techniques (e.g., late blastocyst stages).

-67-

Bovine embryos did not yield to attempts to culture them in vitro past the 8-16 cell "block" until Camous et al. (1984) J. Reprod. Fert. 72, 479-485 demonstrated cleavage to 216 cells when embryos were co-cultured with  
5 trophoblastic tissue.

The co-culture procedure was extended to oviductal tissue, based on the ability of homo- or hetero-oviducts to support development from zygote to blastocyst. Thus, bovine embryos co-cultured with oviductal tissue, or in  
10 medium conditioned by oviductal tissue, developed from zygote to blastocyst in vitro (Eyestone and First, (1989) J. Reprod. Fert. 85, 715-720; Eyestone W.H. (1989) "Factors affecting the development of early bovine embryos in vivo and in vitro." Ph.D. Thesis,  
15 University of Wisconsin). Blastocysts have been produced in this system after superovulation and artificial insemination, or by in vitro maturation (IVM), and fertilization (IVF) of immature oocytes. Blastocysts produced in this fashion resulted in  
20 pregnancies and live calves after transfer to recipient animals. The results obtained were as follows:

<u>Step</u>	<u>Efficiency (%)</u>	<u>Number (per 100)</u>
IVM	90	90
25 IVF	80	72
IVC	30	22
Embryo transfer (% pregnant)	50	11

Therefore, from an initial daily harvest of 500 oocytes,  
30 it is expected the approximately 55 pregnancies will result.

-66-

Preparation of Oviduct Tissue  
Co-Culture and Conditioned Medium

1. Obtain bovine oviducts after slaughter or by salpingectomy.
- 5        2. Harvest luminal tissue by scraping intact oviduct gently with a glass slide.
3. Wash tissue 5 times in 10 ml modified tyrodes-hepes solution (Parrish et al. (1988) Biol. Reprod. 38, 1171-1180).
- 10       4. Resuspend final tissue pellet in M199 + 10% fetal calf serum at a ratio of 1 volume tissue:50 volumes of media.
5. Tissue suspension can be used for embryo-co-culture.
- 15       6. Alternatively, media may be conditioned for 48h; after centrifuging the suspension, the supernatant may be used as embryo culture medium. Conditioned medium may be stored at -70°C, if desired. Conditioned medium should be used at full strength for embryo
- 20 culture (no dilution) (Eyestone (1989) ibid).

EXAMPLE 7

Microinjection of hLF  
Transgene into Bovine Pronuclei

- The DNA fragment containing the hLF expression unit is
- 25 excised from the vector by digestion with the appropriate restriction enzyme(s) and separated on agarose gels. The fragment is purified by electroelution, phenol and chloroform extraction and ethanol precipitation (Maniatis et al.). The DNA
- 30 fragment is dissolved in and dialyzed in 10 mM tris, 0.1 mM EDTA pH 7.2 at a concentration of 1 to 2 µg/ml. Microinjection needles are filled with the dialyzed DNA solution.

-69-

Before in vitro fertilization, cumulus cells are removed from the egg by either vortexing at maximal speed for 2 minutes or pipetting the eggs up and down several times in a standard micropipet. ~~Bovine pronuclei are~~  
5 ~~injected in principle as murine pronuclei~~ (Hogan, B. et al. (1986) in: Manipulating the mouse embryo, Cold Spring Harbor Laboratory) ~~with an additional centrifugation step in order to visualize the pronuclei~~. The injection takes place 18-24 hours after  
10 fertilization. The time varies depending on the bull used as a source of semen. Different batches of semen cause the nuclei to become visible at different times.

Bovine oocytes, matured and fertilized in vitro, are spun in an eppendorf tube in 1 ml of tyrodes-hepes  
15 solution (Parrish (1987)) at 14500 g for eight minutes (Wall et al. (1985) Biol. Reprod. 32, 645-651). The embryos are transferred to a drop of tyrodes-hepes solution on a microscope slide covered with paraffin oil. Using a hydraulic system the oocytes are fixed to  
20 the egg holder in such a way that both the pronuclei are visible (using interference-contrast or phase contrast optics). If necessary, the oocytes are rolled to change their position on the egg holder to visualize the pronuclei. The injection needle is brought into the  
25 same sharp focus of one of the pronuclei. The needle is then advanced through the zona pellucida, cytoplasm into the pronucleus. A small volume of 1-3 pl is injected (containing 20-100 DNA copies) into the pronucleus either by using a constant flow or a pulse  
30 flow (using a switch) of DNA solution out of the needle. Alternatively, two cell stage embryos are spun as described and the nuclei of both blastomers are injected as described. The injected embryos are then transferred to a drop of co-culture medium as described in Example 6  
35 in order to develop to the morula or blastocyst stage.

-70-

EXAMPLE 8Early Detection of  
Transgenesis with hLF Transgene

Upon the microinjection of a construct, the oocyte is  
5 cultured. A proper site of each embryo is cleaved and  
subjected to lysis (King, D. et al. (1988) Molecular  
Reproduction and Development 1, 57-62), proteolysis  
(Higuchi, R., (1989) "Amplifications (A forum for PCR  
Users." 2, 1-3) and DPNI digestion. PCR is performed  
10 as described previously (Ninomiya, T. et al. (1979)  
Molecular Reprod. and Devel. 1, 242-248) with sets of  
two primers, one in  $\alpha$ S1 and the other in hLF cDNA  
sequence. For example, in a PCR where the forward  
primer (30mer)  $\alpha$ S1 sequence is

15 ATG AAA CTT ATC CTC ACC TGT CTT GTG  
(Seq. ID No.: 16)

and the reverse primer (30mer) in hLF sequence is GGG TTT  
TCG AGG GTG CCC CCG AGG ATG GAT (Seq. ID No.: 17); 971-  
1000 of Figure 1), a 990 bp fragment will be generated.  
20 This fragment contains the hitherto inactivated DpNI  
site by loss of adenosine-methylation, at 934 bp away  
from the start of the forward primer.

EXAMPLE 9Production of hLF in Milk of Bovine Species

25 Bovine morula developed from microinjected oocytes are  
split according to the method of Donahue (Donahue, S.  
(1986) Genetic Engineering of Animals, ed. J. Warren  
Evans et al., Plenum). One half of the morula is kept  
in culture to develop into blastocysts. The other half  
30 is subjected to the DNA analysis as described in  
Example 8. When the result of this analysis is known,  
the morula kept in culture are developed into a



-71-

blastocyst or as a source for nuclear transfer into enucleated zygotes. Blastocyst transfer into synchronized cows is performed according to the method of Betteridge (Betteridge, K.J. (1977) in: Embryo transfer in farm animals: a review of techniques and applications).

hLF is detected in the milk of lactating transgenic offspring using the RIA of Example 5.

#### EXAMPLE 10

##### 10      Bovine $\alpha$ S1 Casein/hSA Expression Plasmids

Three overlapping phage clones that contain the complete hSA gene are used to construct an expression vector for hSA. They are designated  $\lambda$ HAL-HA1,  $\lambda$ HAL-3W and  $\lambda$ HAL-H14. They are described in Urano, et al. (1986), J. Biol. Chem., 261, 3244-3251; and Urano, et al. (1984), Gene, 32, 255-261. The sequence of the gene plus some surrounding regions is published in Minghetti, et al. (1986), J. Biol. Chem., 261, 6747-6757. A single phage containing the complete hSA gene is constructed as follows:

Clone HA-1 is cut with BstEII and AhaII. The  $\approx$ 1400 bp fragment running from position 1784 (in the first exon, just downstream of the ATG) to 3181 is isolated and a synthetic linker is attached to the BstEII site at the 5' end containing the first few amino acids that are cut off with BstEII as well as the sequence surrounding the ATG as well as a few convenient restriction sites. This fragment is called fragment #1.

Clone 3W is cut with AhaII and SacI the  $\approx$ 13.1 kb fragment running from position 3181 to 16322 is isolated and a synthetic linker is attached to the SacI site to

-72-

facilitate cloning in phage EMBL3. This fragment is called fragment #2.

These two fragments are ligated and cloned in phage EMBL3. After identification of the correct phage, a  
5 fragment running from just upstream of the BstEII site (where unique restriction sites have been introduced) to the SacI site are isolated and ligated from a SacI to SalI fragment (running from position 16322 to  $\approx$ 21200 isolated from clone H-14. These two fragments are then  
10 ligated and cloned in EMBL4.

After cutting with ClaI (just upstream of the BstEII site, newly introduced) and BamHI (just downstream of the SalI site in the phage DNA) this new clone yields a fragment containing the complete hSA gene with about  
15 2.5 kb 3'-flanking sequence.

To construct an expression vector for hSA cosmid cGP1HLF is partially digested with ClaI and BamHI. This removes the signal sequence, the coding sequence of hLF, the 3'-UTR and poly(A) addition region of  $\alpha$ S1-casein as well  
20 as a small region 3' of the casein gene.

This is ligated to the hSA fragment described above and the resulting cosmid is called cGP1HSA.

The expression vector so formed contains, (1) 16 kb of promoter sequences derived from the  $\alpha$ S1-casein gene,  
25 (2) the first exon and intervening sequence of this gene both present in GP1, (3) the signal sequence of the hSA gene the complete genomic gene coding for hSA including 2.5 kb downstream of that gene, and (4)  $\approx$ 8 kb of 3'-flanking sequence derived from the  $\alpha$ S1-casein gene.

30 This transgene is used to produce transgenic bovine species producing hSA in their milk in a manner

-73-

analogous to that used to produce hLF in the milk of bovine species.

#### EXAMPLE 11

##### Purification of HSA from the Milk of Bovine Species

5

Purification of heterologous proteins from milk is facilitated by the fact that, following casein precipitation, those proteins, for the most part, are found in the whey fraction which is less contaminated than the production media used in microbial or cell-based systems.

10

Chromatographic techniques are preferred for the purification of hSA from cow milk. This approach produces a better recovery and higher albumin purity as well as a lower content of albumin polymers as compared with ethanol fractionation (Curling (1980) in: "Methods of Plasma Protein Fractionation", Curling, ed., Academic Press London, UK; Curling et al. (1982) J. Parenteral Sci. Technol. 36, 59; Berglof et al. and Martinache et al. (1982) Joint Meeting IHS-ISBT, Budapest). The specific transport role of hSA as well as its major role in maintaining intravascular osmotic pressure may also be better preserved upon chromatographic purification (Steinbruch (1982), Joint Meeting ISH-ISBT, Budapest).

15

20

The following steps are used to recover hSA produced in the milk of transgenic cows:

1. Precipitation of caseins (about 80% of milk protein) and essentially all the milk fat at pH 4.5 and/or by adding chymosin. The whey fraction contains the albumin;

30

-74-

2. Affinity-chromatography of albumin on Cibacron blue 3GA-Sepharose CL-6B (Harvey (1980) in: Methods of Plasma Protein Fractionation, op. cit.) This step serves both to remove proteins other than albumin and to decrease the volume to be handled about 30-fold. Albumin is eluted from this matrix with 0.15 M NaCl and 20 mM sodium salicylate at pH 7.5;
  3. Buffer-exchange on Sephadex G-25: desalting into 0.025 M sodium acetate, adjustment to pH 5.2, followed by filtration;
  4. Anion-exchange chromatography on DEAE-Sepharose CL-6B. Desorption of albumin at pH 4.5;
  5. Cation-exchange chromatography on CM-Sepharose CL-6B. Albumin elution with 0.11 M sodium acetate, pH 5.5 and concentration of albumin at a 6% (w/v) solution by ultrafiltration; and
  6. Gel filtration on Sephacryl S-200. Fraction of high-molecular weight protein (e.g. albumin polymers, pyrogens) is discarded. The main fraction (albumin monomers) is concentrated by ultrafiltration and formulated.
- It is to be noted that steps 3-6 are essentially identical to the method described by Curling and others (Curling (1980) op. cit.; Curling et al. (1982) op. cit.; Berglöff et al. (1982) op. cit.) for the purification of hSA from plasma.

#### EXAMPLE 12

Transgenic Mice Containing the Human Serum Albumin (hSA) Transgene  
Generated by Homologous Recombination

Three overlapping genomic hSA clones were used to generate the hSA gene in transgenic mice,  $\lambda$ HAL-HA1,  $\lambda$ HAL-H14 and  $\lambda$ HAL-3W, are shown in Figure 8 as reported

-75-

by Urano, et al. (1984), Gene, 32, 255-261 and Urano, et al. (1986), J. Biol. Chem., 261 3244-3251. Briefly, a genomic library was constructed from a partial EcoRI digest of human fibroblast DNA. For the clones  $\lambda$ HAL-H14  
5 and  $\lambda$ HAL-3W, this library was screened with  $^{32}\text{P}$ -labeled human albumin genomic clones by hybridization in 1 M NaCl, 50 mM Tris-HCl (pH 8.0), 10 mM EDTA, 0.1% SDS, 100 ug/ml of sheared salmon sperm DNA and 10x Denhardt's solution at 65° C overnight after prehybridization in  
10 3x SSC and 10x Denhardt's solution. Following hybridization, filters were washed in 0.2x SSC and 0.1% SDS at 65° C. The isolation of the  $\lambda$ HAL-HA1 clone was identical except that a 0.9 kb BglIII-EcoRI fragment from the 5' end of  $\lambda$ HAL-3W was used to screen the human  
15 fibroblast library.

These three hSA phage clones were used to generate three overlapping linear DNA fragments, which in composite comprised the whole HSA gene and flanking regions. The 5' most fragment I was a EcoRI-EcoRI fragment isolated  
20 from  $\lambda$ HAL-HA1; the middle fragment II was a AclI (=AhaII) -SacI fragment of  $\lambda$ HAL-3W; and the 3' most fragment III was a XhoI-SalI fragment of  $\lambda$ HAL-H14 (Fig. 7). The fragments were treated with klenow DNA polymerase and dNTP's to fill in overhanging sticky  
25 ends. In some experiments, the blunt ended fragments were then treated with bacterial alkaline phosphatase to remove the 5' phosphate groups from each fragment. The overlapping DNA fragments were next concentrated then coinjected into the male pronuclei of fertilized  
30 mouse eggs according to published methods (Hogan, et al. (1986) in "Manipulating the Mouse Embryo: A Laboratory Manual", Cold Spring Harbor Laboratory). While the number of molecules injected varied from  $\approx 25$  to  $\approx 100$  per egg cell, the ratio of the individual fragments was  
35 approximately 1:1:1. Embryos were implanted into the

-76-

uteri of pseudo pregnant female mice according to the methods of Hogan, et al., supra.

To assay correct homologous recombination of the three overlapping fragments and integration of the nascent transgene into the mouse genome, genomic DNA from the newborn pups was subject to the following specific digestions followed by Southern hybridization with HSA cDNA probes:

Bst EII: cuts outside the HSA gene region and yields an 18 kb band if correct recombination occurred;

Nco I: cuts outside the overlapping regions and yields bands of 8.0 and 9.3 kb if correct recombination occurred;

Nco I + Hind III: cuts at several positions outside the region of overlap, indicative of the presence of intact fragments;

Hinc II: cuts in the overlapping regions, yielding several bands indicative of correct arrangement in these regions.

In an initial experiment of 28 transgenic animals born, 22 had correctly recombined all three fragments. From 20 out of those 22 animals blood was collected and assayed for the presence of hSA protein using a radio immuno assay. 15 out of those 20 animals showed hSA expression at levels between 0.5 and 5  $\mu\text{g/mL}$ . None of the animals that had no recombination or that were not transgenic showed any expression. Using RNA blots, only two (the two with the highest protein level) showed a band. We are currently performing blots on RNA that has been enriched for the presence of mRNA (i.e., poly(A)+ RNA). Using reverse transcriptase to synthesize cDNA,

-77-

followed by PCR, we have observed a perfect relationship between the presence of RNA and protein. However, in this experiment we could not determine the size(s) of the RNA.

5

EXAMPLE 13Alternate Construction  
of Transgenes Encoding hLF

This example describes the construction of two hLF transgenes wherein the first contains approximately  
10 16 kb of  $\alpha$ S1 casein 5' expression regulation sequence (pGP1hLF (16 kb) also referred to as p16,8HLF4) and the second contains approximately 7.9 kb of  $\alpha$ S1 casein 5' expression regulation sequence (pGP1hLF (8 kb) also referred to as p8.8HLF4). The overall strategy for  
15 these constructions is depicted in Fig. 9.

A 1.8 kb EcoRI-BglII fragment (fragment C in Fig. 9) was isolated from phage clone GP1. This fragment runs from position -100 of the transcription start site into the second exon of the  $\alpha$ S1 casein gene. The BglII site lies  
20 at the junction of the first intron and second exon of the  $\alpha$ S1 casein gene. The 3' end containing the BglII site was ligated to a synthetic BglII-ClaI linker and subcloned into the plasmid pUC19. The resulting plasmid is designated pEBS.

25 Fragment B in Fig. 9 was isolated as an EcoRI fragment and cloned into the EcoRI site of pEBS. Fragment B includes sequences from position -7500 to position -100 of the transcription start site in the  $\alpha$ S1 casein gene. The plasmid so formed is designated pEB3S and contains  
30 the combination of fragments B and C is the 8.9 kb EcoRI-ClaI fragment running from position -7500 to position +1400 of the transcription start site. The

-78-

8.9 kb EcoRI-ClaI fragment from pEB3, obtained by complete digestion with ClaI and partial digestion with EcoRI was isolated and subcloned into EcoRI-ClaI cut pKUN2 (a derivative of pKUN; Gene (1986) 46, 269-276 containing a NotI restriction site) to form pNE3BS.

An 8.5 kb ClaI-EcoRI fragment (fragment A in Fig. 9) running from position -16000 to position -7500 of the transcription start site was isolated from phage GP1. It was thereafter subcloned into pUC19 to form pSE. Using synthetic oligonucleotide, a unique NotI site was introduced into the ClaI site thereby destroying it. The resulting plasmid is designated pNE.

The insert from pNE was isolated as a NotI-EcoRI fragment and together with the EcoRI-ClaI insert from pNE3BS was ligated into the cloning vector pKUN2. The resulting plasmid pGP1 ( $\Delta$ 2ex) contains 16 kb of  $\alpha$ S1 casein promoter plus the 5' end of the gene to the BglII site at the border of the second exon.

The final plasmid (16,8HLF4) containing the transgene was assembled using the NotI-ClaI fragment from clone pGP1 ( $\Delta$ 2ex) and the Xho-NotI fragment from clone pHLF 3' 10 kb. The structure of this transgene is the same as previously described herein.

As a minor modification to this plasmid the SalI site of this plasmid was removed by cutting with SalI and inserting a linker that contains a NotI site, but not a SalI site. Subsequently, a SalI site was introduced just downstream of the hLF sequence by cutting the KpnI site as that position adding the following linker:

5'-CGTCGACAGTAC-3' (Seq. ID No.: 18)  
CATGGCAGCTGT-5' (SEQ. id no.: 19)



-79-

In effect, the hLF sequence is now surrounded by two unique restriction sites (ClaI and SalI) and can be replaced by any recombinant ANA sequence that has a ClaI-site at the 5'- end and a SalI-site at the 3'- end.

- 5 Another transgene was constructed that is identical to the foregoing except that it contains only about 8 kb of 5'  $\alpha$ S1 casein expression regulation sequence. It was constructed by taking the NotI-ClaI fragment from pNE3BS and fusing it directly into Xho-otI fragment from clone  
10 pHLF 3' 10 kb. The resulting plasmid was designated pGPIhLF (7 kb) (also referred to as p8.8HLF4).

- Plasmid 16,8hLF4 was modified to contain a hybrid splice signal ( $\alpha$ S1 casein-IgG) described in examples 3 and 5. The resulting plasmid was designated 16,8hLF3 and is  
15 identical to 16,8hLF4 except for the presence of a hybrid intron versus a "natural" casein intron in the 5'-UTR.

- The hLF signal sequence can also be used in all of the cDNA constructs disclosed herein instead of the casein  
20 signal sequence. This can be done in the following way: A synthetic oligo was made that contains the complete hLF signal sequence (see Fig. 2) plus a ClaI restriction site at the 5'-end and an EagI restriction site at the 3'-end. These restriction sites also border the casein-  
25 signal sequence in the other plasmids (e.g., p16,8hLF4). A fragment containing the hLF-cDNA surrounded by ClaI and SalI sites was cloned in pGEM7 (Stratagene, Inc.) containing a ClaI and SalI site. The resulting plasmid was digested with ClaI and EagI and used as a vector to  
30 accommodate the ClaI-EagI fragment containing the hLF sequence. From the positive clones, the cDNA, with its own sequence, was excised as a ClaI-SalI fragment and inserted in ClaI-SalI digested p16,8hLF4 to generate p16,8hLF5. Similarly, this Cla-Sal fragment containing

-80-

the hLF-cDNA plus hLF signal sequence can be inserted in any hLF cDNA vector.

#### EXAMPLE 14

##### 5     Production of Recombinant Human Lactoferrin and Human Serum Albumin in the Milk of Transgenic Mice

Transgenic mice were generated utilizing several of the transgenes identified in the examples herein. The transgenes used are identified in Tables 3 and 4. In each case, the 5' and 3' expression regulation sequences  
10 were from the bovine  $\alpha$ S1 casein gene, the RNA splice signal in the 5' untranslated region was either homologous from the  $\alpha$ S1 casein gene or a hybrid casein-IgG intervening sequence. The recombinant DNA in each case was derived from cDNA clones.

15 The transgene containing 26 kb of 5'  $\alpha$ S1 casein expression regulation sequence was generated by in vivo homologous recombination of overlapping fragments. Briefly, a phage clone containing an approximately 14 kb SalI insert was identified. This insert contains about  
20 11.5 kb of sequence upstream from the 3' casein sequence contained in 16,8hLF4 and about 2.5 kb of overlapping sequence. The NotI insert from 16,8hLF4 and the SalI phage insert were coinjected to produce the 26.8hLF4 mice.

-81-

TABLE 3\*

<u>Plasmid from which transgene excised</u>	<u>Length of 5'- expression regulator segment (kb)</u>	<u>Length of 3'- expression regulator segment</u>	<u>IVS</u>	<u>Number of strains</u>	<u>Maximum expression levels recorded (<math>\mu</math>g/ml)</u>	<u>Range of mean levels recorded (<math>\mu</math>g/ml)</u>
p0.7,8 hLF4	0.68	8	homologous	6	0.0-0.8	0.0-0.1
p8,8 hLF4	6.2	8	homologous	6	5-36	2.5-16
‡p16,8 hLF4	14.5	8	homologous	5	0.3-3.6	0.0-1.8
p26,8 hLF4	26	8	homologous	5	0.6-10	0.2-1.7
p16,8 hLF3	14.5	8	heterologous	13	0.0-708	0.0-200

\*The number in the plasmid designations before the comma represents the approximate length in kbp in the 5' sequence from the bovine  $\alpha$ S1 casein promoter/flanking region while the number after the comma represents the approximate length in kbp in the 3' flanking sequence of the  $\alpha$ S1 gene. Note the actual number of bases for the 8 kb and 16 kb promoter (5' flanking region) is 6.2 and 14.5 kbp, respectively.

‡Exception: An additional p16,8hLF4 transgenic mouse (line 145) not included in the data in Table 3 gave a maximum expression level of 224  $\mu$ g/ml and a mean of 112  $\mu$ g/ml.

-82-

TABLE 4

16,8hLF 3 Expression Data5/13 - High Expressors:

	Mouse line No.	Mean level of expression range ( $\mu$ g/ml)	Maximum level of expression range ( $\mu$ g/ml)
5	27	33.5	97.5
	29	37.5	66.0
	32	21.2	148.0
10	33	200.0	708.0
	38	25.0	126.0

8/13 - Low Expressors:

0.0-1.7

0.2-18

The data in Tables 3 and 4 demonstrates that the hybrid  
 15 intron + heterologous splice acceptor site dramatically  
 increases expression levels in a significant number of  
 cases (5/13).

The construct 16,8hLF4 is expressed at high level (in  
 same range as 16,8hLF3). However, (in mice) this only  
 20 occurs in a small number of cases and 1/16 when 8,8hLF4  
 and 26,8hLF4 are included). Similar results were  
 obtained using a hSA cDNA.

Briefly, the 16,8hSA4 transgene was constructed by  
 digesting p16,8hLF4 with ClaI and SalI to remove the hLF  
 25 cDNA sequence. hSA cDNA was excised from a clone with  
 EcoRI. A ClaI synthetic linker was added to the 5'  
 (upstream) end and a SalI linker to the 3' (downstream)  
 end. After insertion into the ClaI/SalI digested  
 16,8hLF4 vector, 16,8hSA4 was formed from which the NotI  
 30 insert was excised and used for microinjection.

The 16,8hSA4 construct yield 9 lines. One of the 9  
 lines gave high level expression (100  $\mu$ g/ml), while the

-83-

remaining 8 of 9 gave low expression (0.01-0.05  $\mu\text{g/ml}$ ). This indicates that the level and the frequency of hLF expression in the mouse mammary gland are not determined by the particular cDNA used, but are an inherent characteristic of the 16,8X4 construct (i.e., the 16 kb 5' and 8 kb 3' flanking regions of the  $\alpha$ -S1 casein gene combined with the heterologous IVS).

The data also show that 0.7 kb of 5'  $\alpha$ -S1 casein flanking sequence does not drive high level expression and that 8 (6.2), 16 (14.5) and 26 kb are more effective. In this respect, 8 kb is slightly more effective than 16 or 26 kb of 5' flanking sequence.

Also, RNA analysis has shown that expression of the cDNA constructs is tissue-specific and stage-specific (i.e., expression is only observed in the lactating mammary gland), that the transcripts are correctly sized and that RNA and protein levels correlate.

#### EXAMPLE 15

##### Generation of hLF Transgenic Cattle

Transgenesis in the bovine system was obtained utilizing the p16,8hLF4 transgene described in Example 13.

##### Oocyte Maturation and Fertilization

Bovine oocytes were collected by aspiration of follicles present on ovaries obtained from slaughterhouses and transported in an insulated container at 30-32°C. Oocytes, together with follicular fluid, were aspirated from 2-8 mm diameter follicles and pooled into 50 ml conical tubes. Cumulus-oocyte complexes (COC) were allowed to settle into a pellet, after which the supernatant was discarded and the pellet washed in 50 ml TL-Hepes (Vander Shaws, et al. (1991) Theriogenology

-84-

35, 288 (Abstr.). COC, containing several intact, unexpanded cumulus cell layers, were selected and isolated under a dissecting microscope at 15 x magnification, washed four times in 10 ml TL-Hepes, once  
5 in 2-3 ml TCM199+10% fetal calf serum (M199) and then paraffin oil (20 COC/droplet). COC were incubated for 23 h in a humidified atmosphere of 5% CO<sub>2</sub> in air at 39°C.

A total of about 2500 oocytes were used. On average, two aspiration sessions occurred per week. The yield  
10 of aspirated oocytes was highly variable from day to day, with a mean daily number of about 150. Maturation and fertilization were analyzed by cytological analysis. Maturation was defined as the breakdown of the nuclear membrane, the appearance of the first polar body and a  
15 metaphase plate. Oocytes were fertilized in vitro with frozen thawed-sperm obtained from three different bulls with excellent characteristics with respect to genetic background, field performance and ease of calving. Sperm capacitation was facilitated with heparin.  
20 Parrish, J. et al. (1986) Theriogenology 25:591-600. Since sperm from individual bulls respond differently to specific fertilization conditions, semen from each lot was tested in advance to determine optimal heparin and sperm concentration required to maximize normal  
25 fertilization frequency and to minimize polyspermy. Fertilization conditions for a given bull were selected after screening at heparin concentrations of 0.0, 1.0 and 10.0 mg heparin/ml, and at 1.0, 2.0 and 4.0 x 10<sup>6</sup> motile sperm/ml. Since the proportion of sperm that  
30 survives freezing and thawing varies from bull to bull (approximately 30-60% for the bulls was used here) sperm preparations were enriched for live, motile sperm by a "swim-up" procedure (Parrish, J. et al. Ibid), alternatively, sperm were centrifuged through a percoll  
35 gradient. After isolation of the motile portion, sperm were counted on a hemocytometer, diluted to an

-85-

appropriate concentration to yield a 25-fold concentrated stock. The fertilization medium consisted of TALP medium (Banister, Bethal Biol. Reprod. 28:235-247) supplemented with 2.0-10.0 mg/ml heparin (from porcine intestinal mucosa, 177 IU/mg; Sigma) and if the cumulus was removed prior to fertilization, 1mM hypotaurine, 10mM penicillamine, 20mM epinephrine and 2mM sodium metabisulfite. Matured COC were selected on the basis of expanded cumulus masses for fertilization, washed once in 10 ml fertilization medium, and either added directly to fertilization droplets, or first stripped of their cumulus investment by gentle pipetting through a small-bore, fire-polished pipet and then added to the droplets. Finally, sperm cells were added to a final concentration of  $1 \times 10^6$ - $2.0 \times 10^6$ /ml. After 16-24 h, presumptive zygotes were removed from fertilization droplets. At this point, 20-30 zygotes for each experiment were fixed in 3:1 ethanol:acetic acid for 24 h, stained with 1% aceto-orcein (in 40% acetic acid), and examined to determine fertilization frequency (percentage of sample with 2 pronuclei and a sperm tail). For each batch of semen, the 'in vitro' fertilization conditions (heparin concentration and sperm number) were optimized to obtain normal fertilization rates ranging from 50 to 70% as determined by the presence of two pronuclei and a sperm tail as described above. Either one of two techniques were used for selection of motile sperm: the swim-up technique and centrifugation through a Percoll gradient. No significant differences in fertilization rates between these methods were recorded. The efficiencies of these and the following steps are shown in Table 5. The remaining oocyte were then prepared for microinjection.

TABLE 5

Efficiencies of the steps involved in the process  
from immature bovine oocytes to transgenic calves

-86-

Step	Total No.	Percent*
oocytes	2470	-
matured	2297	93
fertilized	1358	61
5 injected	1154	85
survival	981	85
cleavage	687	70
transferred	129 <sup>†</sup>	1.9
pregnant	21	2.1
10 integration	2	1.0
*Percentages indicate the proportion of embryos or cells that successfully complete each step. <sup>†</sup> Sixty-nine transfers of single blastocytes resulting in 7 pregnancies; 30 transfers of twinned embryos, resulting in 14 pregnancies.		

Microinjection. The 26 kbp casein-hLF transgene (from p 16,8hLF4) used for microinjection was released by NotI digestion and purified by agarose gel electrophoresis and electroelution. The final DNA concentration was adjusted to 2.5  $\mu$ g/ml. Batches of 50 cumulus-intact fertilized oocytes were stripped either as described above or by vortexing 2 minutes in 2ml TL-hepes medium in a 10ml conical tube. In order to visualize the pronuclei, cumulus free oocytes were centrifuged in 1ml TL-hepes medium 8 minutes at 14,500 x g in an Eppendorf centrifuge. Wall, R. et al. Biol. Reprod. 32:645-651. ~~Microinjection was performed essentially as described by Hogan B. et al. (1985) Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y..~~

Embryo culture. Embryos were cultured from the zygote to the compact morula or blastocyst stage in oviductal-tissue conditioned medium. Eyestone et al., (1991) J. Reprod. Fert. 92:59-64. Oviducts were obtained at slaughter and transported at ambient temperature. Luminal tissue from 2-4 oviducts (1-2 cows) was harvested by gently scraping intact oviducts on the outside with a glass slide. The extruded material was



-87-

washed 5 times in 10 ml TALP-Hepes and diluted in M199 to a tissue:media ratio of 1:50. Media were conditioned in 50 ml "T" flasks containing five ml of oviduct tissue suspension. Conditioned media frequently contained a proteinaceous precipitate after thawing, which was removed by centrifugation. Droplets were covered with paraffin oil and were incubated for 2 h to permit pH to equilibrate prior to adding zygotes. Zygotes were placed in culture droplets within 2 h after microinjection. Initial cleavage (>2 cells) was assessed 42 h after adding sperm. Media were not changed during the course of incubations. Criteria for normal development consisted of attainment of the compact morula or blastocyst stage.

**Embryo transfer.** The synchronization schedule was set up so that recipients started estrous on the same day at which oocytes were aspirated from slaughterhouse ovaries (i.e., start of maturation is day 1). Estrous in recipient cattle was synchronized with a 9-day Norgestamet (Intervet, Boxmeer, The Netherlands) treatment (administered in an ear implant according to the manufacturer), and a 500 µg dose of cloprostanol given on day 7 of the Norgestamet treatment. Estrous occurred within 2-3 days after implant removal. Embryos were transferred non-surgically to recipient heifers 5-7 days after estrous (1-2 embryos/uterine horn). Recipients received 9-day old embryos, at which time they have developed to the compact morula or early blastocyst stage. These embryos are one day ahead in development compared to the stage of the estrous cycle of the recipients. In case of two microinjection sessions on subsequent days, one group of recipients was used that were in synchrony with the first batch of oocytes collected. Transfers of embryos that developed from oocytes aspirated on the day of the start of estrous gave better results than embryos from oocytes

-88-

obtained one day later. Due to the somewhat delayed development of microinjected embryos, there appeared to be a better synchrony between the recipients and the first group of embryos. Recipients received two embryos when the quality grade (according to Linder and Wright, Theriogenology 20:407-416) was fair to poor and only one single embryo when the quality grade was excellent to good. Each pregnant recipient that received 2 embryos carried only one fetus to term. The overall pregnancy rate was 21%, which is significantly less than the rates reported by others with non-microinjected embryos which had developed in vivo (Linder and Wright, Ibid and Massy et al. (1984) Theriogenology 21:196-217). In the experiments described here, no transfers with non-injected embryos were performed.

Pregnancy was determined by rectal palpation at 45 to 60 days of gestation. A total of 21 pregnancies were established (confirmed by rectal palpation 45-460 days after transfer). During pregnancy, 2 fetuses were lost. One recipient aborted spontaneously for unknown reasons at 7.5 months of gestation. The second fetus, collected at slaughter of the recipient at 3 weeks after the calculated day of parturition, was a full grown dead calf having an abnormal embryonic development called 'schistosoma reflexum'. In both cases, no intact DNA could be isolated for analysis. Nineteen calves were born after normal pregnancies. One of these calves died during parturition, and a second, 24 hours after birth, because of pneumonia following accidental inhalation of milk. A third calf, born after a pregnancy of 10 months and with a body weight of 70kg was euthanized at an age of 3 weeks. Pathological analysis indicated that the animal was suffering from sepsis due to chronic omphalophlebitis. Tissues that could be analyzed from the three dead calves contained no integrated human lactoferrin (hLF) sequences. Therefore, the cause of

-89-

their death is unlikely to be related to transgene integration. The remaining 16 calves are in excellent health.

**Structure of the transgene.** In Fig. 12A, the coding  
5 sequence of the hLF cDNA is depicted by a hatched box.  
The position of the translational start and stop codon  
is indicated. The 5' and 3' untranslated regions are  
encoded by  $\alpha$ S1 casein exons (open boxes). Intervening  
10 sequences interrupting these exons are represented by  
a single line. The expression unit is surrounded by  
flanking sequences derived from the bovine  $\alpha$ S1 casein  
gene (indicated by a double line). Positions of  
restriction enzyme sites are indicated by the following  
symbols: R, EcoRI; A, Asp718; N, NotI. The NotI sites  
15 are not present at the indicated positions in the bovine  
 $\alpha$ S1 casein gene itself, but were introduced by synthetic  
linkers. The black bar represents the position of the  
probe used to detect the presence of the transgene.  
Sizes of the fragments (in kbp) obtained after digestion  
20 with EcoRI or Asp718 are shown at the bottom.

**DNA analysis.** DNA was isolated from placenta, blood and  
ear tissue from all calves. A Southern blot analysis  
of the extracted DNA is shown in Fig. 12B. Ten  $\mu$ g of  
DNA was loaded per lane. Fragment size markers are in  
25 kbp (HindIII digest of lamda DNA) are indicated on the  
left. Lane 1, EcoRI digested human DNA (isolated from  
blood), lane 2, Eco RI digested DNA from calf #4  
isolated from blood; lane 3, Asp718 digested DNA from  
calf #4 isolated from blood; lane 4, EcoRI digested  
30 placental DNA from calf #4; lane 5, Asp718 digested  
placental DNA from calf #4; lane 6, EcoRI digested DNA  
from calf #15 isolated from blood; lane 7, Asp718  
digested DNA from calf #15 isolated from blood; lane 8,  
EcoRI digested DNA from calf #15 isolated from ear  
35 tissue; lane 9, Asp718 digested DNA from calf #15

-90-

isolated from ear tissue; lane 10, EcoRI digested placental DNA from calf #15; lane 11, Asp718 digested placental DNA from calf #15; lane 12, EcoRI digested DNA isolated from the tail of a transgenic mouse harboring the same construct. DNA extraction, Southern blot analysis and hybridization were performed according to standard procedures. The probe used in the Southern blotting experiment was a 758 bp EcoRV-EcoRI fragment covering the 3' part of the hLF cDNA of Figure 2.

10 Southern blot analysis using hLF cDNA as a probe indicated that in tissues of two calves (#4 and #15) transgene sequences had been integrated into the host genome. Calf #15 (a female) was mosaic for integration of the transgene. Placental tissue was positive, whereas in blood and ear tissue no hLF sequences could be detected. The copy number in the placenta was 1-2. The restriction enzyme map of the transgene was different from that expected based on the map of the casein-hLF plasmid (FIG. 12A) and based on the pattern obtained in many individual transgenic mice (data not shown). Apparently, a rearrangement had occurred involving a deletion of part of the DNA construct. It is not clear whether this rearrangement event is related to the fact that the transgene could not be detected in all tissues. In mice, it has been shown that over 30% of all transgenic animals born are mosaic.

Calf #4 (a male) showed, in all three tissues, the same hybridization pattern that was identical to the expected one. Restriction digestions with different enzymes indicated that head-to-tail concatamers of intact copies had integrated and there was no indication of rearrangements. Copy numbers were estimated by comparing the intensities of the transgenic band with bands resulting from hybridization of the hLF probe to human DNA (FIG. 12B). In calf #4, between 5 and 10

-91-

copies of the transgene had integrated in all three tissues examined.

An analysis of sperm produced by calf #4 detected no abnormalities. DNA was subsequently isolated from the sperm and analyzed for the presence of the hLF-transgene. It appeared that the copy number of the transgene (2-3) was the same in sperm as in other tissues indicating that calf #4 is not mosaic and should be able to transmit the transgene to 50% of his offspring.

#### EXAMPLE 16

##### Construction of Transgene Cassette for Genomic Recombinant DNA

The plasmids described so far all contain regions derived from the bovine  $\alpha$ S1-casein untranscribed regions (including intervening sequences). When a genomic gene is to be expressed that already contains untranslated regions and intervening sequences permissive for high expression, it is preferable to use expression cassettes where the flanking regions including the transcription initiation site of the  $\alpha$ S1 casein gene are operably linked to the untranslated regions of the gene to be expressed. Such an expression cassette is p-16 kb,CS and was constructed as follows: plasmid pS1 3'5'hLF was used as a template in a PCR experiment. This plasmid contains 680 bp of promoter sequence of the  $\alpha$ S1 casein gene as well as its first exon. the rest of this plasmid is not relevant for this experiment. The upstream primer was located just upstream of the insert in the plasmid moiety (just upstream of a NotI restriction site). Its sequence is: 5'-CGA CGT TGT AAA ACG ACGG-3'.

-92-

The downstream primer was located in exon 1. Its sequence matches the first 19 bp of the exon exactly and also has a non-hybridizing region of 17 bp containing a ClaI and a SalI site. It has the following sequence:

5' -ATTGTCGACTTATCGATGGGTTGATGATCAAGGTGA-3'

The amplified fragment was digested with NotI and SalI and ligated into pKUN2 (see Example 13). The resulting plasmid (p-680CS) therefore harbors a proximal promoter fragment from -680 to +19, plus two restriction sites just downstream of those 19 bp.

This plasmid was digested with NotI (just upstream of -680) and NsiI (at -280) and used as a vector to ligate to a fragment running from a NotI site (just upstream of -16 kb) to NsiI (-280) isolated from p16,8hLF4 (Example 13). This plasmid (p-16 kb,CS) therefore harbors a promoter fragment from  $\approx$ -16,000 to +19. It can be used to insert genomic genes that carry their own UTR's and poly(A)-signal. After insertion of the genomic gene as a ClaI-SalI fragment, the  $\alpha$ S1 casein 3'-flanking region can be inserted as a SalI-fragment.

#### EXAMPLE 17

##### Construction of Transgene for Production of Protein C

The genomic sequence of Protein C has been published. Foster, et al. (1985) Proc. Natl. Acad. Sci. USA 82, 4673-4677. This sequence, however, does not include the first exon which was identified through the cDNA sequence published by Beckman, et al. (1985) Nucl. Acids Res. 13, 5233-5247. The first exon of Protein C is located at position -1499 to -1448 in the Foster sequence. The transgene for expressing and secreting


-93-

Protein C into the milk of bovine species is shown in Fig. 10. This transgene was constructed as follows.

A human genomic library in EMBL-3 (Clonotech) is probed with a sequence specific for protein C. A purified  
 5 phage DNA prep containing the complete Protein C gene is isolated. The phage is isolated from an *E. coli* strain having the Dam phenotype, such a strain GM113. This results in cloned DNA which is not methylated and as such all *Cla*I restriction sites can be cleaved.

10 A *Cla*I *Nhe*I fragment running from positions +1333 to 11483 is isolated. This is designated fragment I.

pGEM7 (Stratogene, Inc.) is digested with *Sph*I and *Sma*I. The region in between is replaced by the corresponding region of plasmid pKUN (Gene (1986) 46, 269-276). The  
 15 resulting plasmid is designated pGEM7A and has the following restriction map in the relevant region:



20 Two primers are synthesized. Primer GP125 has the following sequence:

5' - CAA ATC GAT TGA ACT TGC AGT ATC TCC ACG AC - 3'  
                   *Cla*I

Primer GP 126 has the following sequence:

25           5' - GGG ATC GAT CAG ATT CTG TCC CCC AT - 3'  
                                   *Cla*I

Primer GP125 has an overlap with exon 0 (position 654 to 675 of the Protein C gene) and introduces a *Cla*I site in the 5' untranslated region. Exon 0 is the exon not

-94-

identified by Foster, et al. Primer GP126 overlaps the region from 1344 to 1315 in the Protein C gene. This region contains a ClaI site.

The region between position 654 and 1344 is amplified  
5 using either human DNA or phage DNA as a template. The so amplified material is digested with ClaI and cloned in vector pGEN7a to form pPCCC. This vector is propagated in a dam negative strain such as GM113 and partially cut with ClaI (only the plasmids that are cut  
10 once with ClaI at position 1340 are of interest) and completely with XbaI. The ClaI NheI fragment (fragment 1) is cloned into this vector. The resultant plasmid is designated pPC. Its structure is shown in Fig. 10. From this plasmid, the Protein C transgene is isolated  
15 as a ClaI-SalI fragment and ligated into p16 kb, CS (See Example 15) to generate a transgene capable of expressing Protein C in bovine milk, this plasmid is designated p16 kb,CS,PC.

The transgene contained within plasmid p 16 kb, CS, PC  
20 is excised with NotI and used to generate transgenic bovine species as previously described. Such transgenic animals are capable of producing protein C in their milk.

#### EXAMPLE 18

25 Human Lactoferrin Transgene Formed by  
In Vivo Homologous Recombination:  
Microinjection of Two Overlapping DNA Fragments

To obtain the entire hLF genomic clone, two human genomic cosmid libraries were screened using an hLF cDNA  
30 clone described herein as a probe. Of 14 clones isolated, 2 clones (designated 13.1 and 13.2; one from each human cosmid library) contained the entire hLF gene



-95-

as determined by hybridization with primers specific for the first and last (17th) hLF exons and by DNA sequencing. The insert sizes of these hLF genomic clones was 42 kbp for clone 13.1 and 43 kbp for clone 13.2. Clones 13.1 and 13.2 contain 5 kbp and 13 kbp of 5' flanking sequences, respectively. The 3' flanking region of clone 13.2 is between 1 kbp and 3 kbp; clone 13.1 contains 7 kbp of additional 3' flanking sequence. The size of the structural hLF gene (= introns + exons) is approximately 30 kb.

The identity of the hLF clones was confirmed by sequencing several exons (incl. first and last) and comparing these sequences and the promoter region to the hLF cDNA sequence shown in Fig. 2. In addition, the clones were transfected into human kidney 293 cells and hLF expression was detected, indicating that both clones were functional.

A comparison of the 13.1 and 13.2 clones (derived from independent libraries) by restriction mapping and Southern blotting revealed no differences in the corresponding regions (i.e. in the structural hLF gene). Southern blotting experiments revealed that the hLF gene is a single copy gene in the human genome. Figs. 13-16 illustrate the overall procedure for generating the  $\alpha$ S1 casein/genomic hLF transgene.

The most 5' ApaI site in the structural hLF gene is located in exon I, in the hLF signal sequence. The 400 bp region immediately 5' of exon I was sequenced. This region contains the transcription initiation site of the hLF gene and a TATA-box. This region also includes a BamHI restriction site.

To construct a mammary gland specific expression vector it was necessary to fuse the 8 (6.2) kbp or 16 (14.5)

-96-

kbp  $\alpha$ S1 bovine casein promoter region to the genomic hLF clone. However, the total size of such a construct, about 50 or 60 kb (6.2 or 14.5 kb from the casein gene promoter + 8 kb from the cosmid vector and 35-40 kb from the hLF genomic clone, i.e., about 50-63 kb), renders the use of conventional cloning vectors difficult. Therefore, the 8 kbp or 16 kbp  $\alpha$ S1 5' casein promoter and flanking sequence was fused to 9 kb of the 5' region of the structural hLF gene (Fig. 15A) and this fragment was coinjected with an overlapping hLF fragment containing about 33 to 34 kbp of the 3' sequence of generic hLF clone 13.1 obtained by ClaI digestion. See Figs. 13 and 16.

The BamHI fragment (containing exon I) from clone 13.2 was subcloned into the plasmid pUC19 (Fig. 14). From this clone, a 8.9 kbp ApaI-SalI fragment was isolated by ApaI (partial digest) and SalI digestion. This fragment lacks most of the hLF signal sequence and all of the hLF 5' UTR. A synthetic sequence (Fig. 15B) representing this missing region was obtained by synthesizing 2 complementary DNA strands (a 68-mer and a 62-mer) which runs from the 5' ApaI site into the downstream region from the hLF TATA-box. After annealing these primers a DNA fragment was generated which has a 5' ClaI overhang and a 3' ApaI overhang. Subsequent sequencing of the cbI-ApCI fragment showed that it has the sequence given in Fig. 15B, which differs at one position from the native sequence. This synthetic ClaI-ApaI fragment and the 8.9 kbp ApaI-SalI fragment described above were ligated into p-16kbCS and into a similar plasmid, containing 8 kbp instead of 16 kbp of the  $\alpha$ S1 casein promoter. This yields two plasmids, containing 16 kbp or 8 kbp of bovine  $\alpha$ S1 casein promoter, fused to the 5' part (9 kbp) of the hLF genomic gene. See Fig. 15A. These fragments were cut out (NotI-SalI) and coinjected with the 3' 33 to 34 kbp

-97-

ClaI fragment from hLF cosmid clone 13.1. The coinjected fragments had an overlap of 5.4 kbp.

Upon coinjection of the constructs containing the 8 kbp  $\alpha$ S1 casein promoter, 8 independent transgenic mice were identified by tail-DNA blotting. To determine if homologous recombination had occurred, chromosomal DNA (from tails of founders and offspring) was digested with ApaI and analyzed by Southern blotting. The 2.7 kb ClaI-MluI fragment (see Fig. 13 or Fig. 17) that is located in the overlap was used as a probe. When homologous recombination occurred, a band of  $7.5 + 0.3 = 7.8$  kb is generated and detected with this probe. This band is also present in human chromosomal DNA, which was used as a control in the analysis. If homologous recombination has not occurred, the probe detects bands of varying size, depending on the location of ApaI sites around the site of integration.

The diagnostic 7.8 kb band was detected in all 8 transgenic mouse lines, indicating that each transgenic mouse contained recombined fragments. For these 8 mouse lines (founder no's: 936, 937, 950, 951, 982, 983, 984 and 985), milk was collected from lactating females (founder and/or offspring) and assayed for hLF protein expression. The data on 7 mouse lines is shown below.

-93-

	<u>Mouse line</u>	<u>Expression Level</u> <u>(max.) (mghLF/ml)</u>
	936	4.5
	937	6.0
5	950	0.003
	951	0.010
	982	5.9
	983	similar to 982 and 937 on day 2 and 4 of lactation*
10	984	2.8
	985	6.6

\*: This mouse died (by accident) on day 4 of lactation. At this time, hLF expression had reached a level of 0.3 mghLF/ml. This is exactly the level found for other high expressors (e.g. lines 937, 982, 984) at this early stage of lactation. This phenomenon of gradual increase of hLF expression at the beginning of, and in particular, the first lactation has been commonly observed by us in the mice generated herein. Therefore, mouse 983 is classified as a high level expressor.

The tissue-specificity of hLF expression was determined by isolating total RNA from a large number of tissues and analyzing for the presence and levels of transgene derived mRNA. Based on this analysis, hLF mRNA only occurs in the lactating mammary gland and expression is tissue- and stage-specific.

RNA levels were below the threshold of detection in lines 950 and 951, but were high in high expressing lines and correlated with bovine  $\alpha$ S1 casein expression levels. This was determined by Northern blot analysis of both bovine lactating mammary gland RNA and mammary gland RNA from lactating transgenic mice. A 24 bp synthetic oligomer which hybridizes to exactly the same

-99-

sequence in the 5' UTR of bovine  $\alpha$ S1 casein RNA and in the transgene RNA was used as a probe. Expression levels were compared directly by quantification of the amount of labelled probe hybridized to the transgene-  
5 and  $\alpha$ S1 RNA. When a correction was made for the size difference between bovine  $\alpha$ S1 casein (20 kD) and hLF (80 kD), the ratio of mRNA to protein was in the same range for bovine  $\alpha$ S1 casein and hLF. This indicates that translation and secretion of the transgene derived hLF  
10 is not impaired. The length of the hLF mRNA was as expected (about 2.5 kb) but in mouse line 937 a longer band (3-3.5 kb) of slightly less intensity was also observed. The occurrence of this band may be related to the homologous recombination process. It remains to  
15 be determined if this RNA translates into bona fine hHL.

It has been suggested that casein promoters are less favorable for obtaining high level expression than other milk specific promoters. However, the present data show that this is not the case. With respect to both  
20 expression level and percentage of expressing animals, the transgenes containing  $\alpha$ S1 casein sequences perform better than any other mammary gland specific transgene reported.

The above data compared to those obtained with  
25 constructs containing hLF cDNA provide the following observation. The best cDNA expression vector herein (16,8hLF3) always expresses at much lower levels as compared to the genomic hLF construct. Of 13 cDNA lines generated, 3 expressed at very low levels (1-5  $\mu$ g/ml),  
30 5 expressed from 40 to 200  $\mu$ g/ml. These relatively low levels (although high for cDNA expression) as compared to that observed for genomic hLF (containing the same flanking sequences) indicate that genomic sequences produce consistently higher expression levels.

-100-

EXAMPLE 19Generation of Genomic Human Lactoferrin Transgenes  
by Conventional Cosmid Ligation Techniques

hLF genomic transgenes have also been generated by  
5 conventional ligations in cosmids. The first construct  
8hLFgen is similar to the transgene generated by  
coinjection, but contains the 3' ClaI fragment from hLF  
clone 13.2. The size of this fragment is about 26-  
27 kb. The second construct 16hLFgen is identical to  
10 8hLFgen, but contains a larger stretch of  $\alpha$ S1 casein  
promoter sequences.

Construction detail:

The NotI-MluI fragment from the construct depicted in  
Fig. 15A (referred to as 8hLFgen9k) was used to prepare  
15 the 8hLFgen construct. This NotI-MluI fragment contains  
the synthetic ClaI-ApaI fragment depicted in Fig. 15B.  
This synthetic sequence contains 24 bp of the hLF 5'-UTR  
and encodes for most of the hLF signal sequence (see  
Fig. 15C). This NotI-MluI was ligated with the 3' MluI-  
20 ClaI fragment from clone 13.2 and a ClaI-NotI linker as  
shown in Fig. 17. The cloning vector was cosmid pWE15  
cut with NotI, from which the internal ClaI and SalI  
sites had been deleted.

The first intron of the hLF gene is located 4 bp  
25 downstream of the ApaI site in the signal sequence. As  
a result, the DNA sequence encoding the 19 aa signal  
sequence is partly located in exon 1 (43 bp, encoding  
14 aa and 1 codon partially) and in exon 2 (the first  
14 bp, encoding 4 aa and 1 codon partially). The exact  
30 position of hLF intron 1 was determined by DNA  
sequencing and comparing the genomic sequence to the hLF  
cDNA sequence. The sequence upstream of the translation  
initiation site (355 bp, containing the hLF 5' UTR and  
5' flanking region) was also sequenced.

-101-

- The hLF transcription initiation site was not included in the genomic hLF constructs as shown. Instead, they contain the bovine  $\alpha$ S1 casein gene transcription initiation site. Although the exact position of the hLF 'cap' site has not been determined, it is probably located about 30 bp downstream of the 'TATA' box, as is the case for the vast majority of eukaryotic genes. In addition, for the mouse LF gene the transcription initiation site has been mapped (Shirsat, et al. (1992) Gene 110, 229-234; Liu and Teng (1991) J. of Biol. Chem. 266, 21880-21885). On the basis of homology between the mLF and hLF 5' UTR, it is concluded that genomic hLF constructs herein do not contain the hLF transcription initiation site.
- 15 The cDNA contains a Thr codon (ACA) at aa position 130 (see Fig. 2). The corresponding region in genomic hLF clones 13.1 and 13.2 (exon 4, plus parts of intron 3 and 4) have been sequenced. These clones contain the sequence ATA, which encodes isoleucine. The cDNA also contains a Cys codon (TGC) at position 404 (see Fig. 2). In hLF clones 13.1 and 13.2 this is a GGC, encoding glycine.

By using the NotI-MluI fragment from 16hLFgen9k instead of from 8hLFgen9k, 16hLFgen was generated.

25 Construction of 8hLFgen37:

- The 5' NotI-MluI fragment from the construct depicted in Fig. 15A (called 8hLFgen9k) was ligated to the 3' MluI-ClaI fragment from clone 13.1, combined with a ClaI-NotI linker (compare Fig. 17: read 13.1 instead of 13.2). The cloning vector was cosmid pWE15, from which the internal ClaI and SalI sites had been deleted, cut with NotI. Prior to microinjection, vector sequences were removed via NotI digestion.

-102-

All constructs were cut from the vector using NotI, and microinjected.

Expression data:

Three mice containing 8hLFgen and 5 mice containing 16hLFgen were generated. Preliminary expression data in milk are as follows:

<u>Construct</u>	<u>Line</u>	Max. hLF expression <u>in milk (mg/ml)</u>
8hLFgen	1089	0.95
	1252	1.2
	1401	1.4
16 hLFgen	1112	2.8
	1113	ND
	1134	0.3
	1185	ND
	1191	ND
8hLF37	1507	4.1
	1556	8.7

ND = not done

SUBSTITUTE SHEET



-103-

EXAMPLE 20Bovine  $\beta$ LG/Human Lactoferrin Transgenes

The bovine  $\beta$ LG-promoter (beta-lactoglobulin) was used to construct a transgene encoding for the expression of hLF. Briefly, the  $\alpha$ S1 promoter in the genomic hLF constructs 8hLFgen and 8hLFgen37 were replaced with the bovine  $\beta$ LG-promoter. The resulting constructs are referred to as  $\beta$ LG-hLFgen and  $\beta$ LG-hLFgen37. The overall strategy for these constructions are depicted in Figs. 18-20.

Isolation of the bovine  $\beta$ LG-promoter.

The charon 28 phage clone  $\lambda$  $\beta$ LG-13, described by Silva et al., (1990) Nucl. Acids Res. 18:3051, was obtained from Dr. Carl A. Batt. This clone was isolated from a bovine genomic library by screening with a  $\beta$ LG cDNA probe. It contains the structural  $\beta$ LG gene and about 8 kb of 5' flanking region. From this clone, a 4.3 kb EcoRI fragment was isolated and subcloned into plasmid pKUN5 using standard procedures (see Fig. 18).

From this plasmid, a 3.2 kb NotI-SacI fragment was isolated. The NotI site was derived from the polylinker of the cloning vector. The SacI site lies 15 bp downstream of the BLG transcription initiation site. A PvuII site is located five bp upstream of the translation initiation site. A fragment representing the region between the SacI and PvuII sites (including these sites) was generated by synthesizing and annealing the 30-mer and 37-mer DNA oligomers depicted in Fig. 18. This fragment also contains a ClaI and a SalI site directly downstream of the PvuII site (Fig. 18). The 3.2 kb NotI-SacI fragment and the synthetic SacI-SalI fragment were ligated into a pKUN plasmid (pKUNdeltaC), from which the internal ClaI site had previously been

-104-

removed by cutting with ClaI and subsequent treatment of the cut vector with Klenow enzyme. This ligation resulted in plasmid pBLG3.2.

5 The 734 bp region directly upstream of the translation initiation site was sequenced and compared to the corresponding region of the published sequence of the sheep BLG promoter (see Fig. 24). Overall homology was 91%, indicating that the sheep- and bovine BLG-promoters are very similar.

10 Generation of  $\beta$ LG-hLF constructs:

The 8.9 ClaI-SalI fragment from construct 8hLFgen9k (Example 15A) was isolated and cloned into p $\beta$ LG3.2 after cutting this vector with ClaI and SalI. This ligation resulted in construct p $\beta$ LGhLFgen9k (Fig. 19). From this  
15 construct the 9.4 kb NotI-MluI fragment was isolated and, together with the 23-24 kb MluI-NotI fragment isolated from 8hLFgen, ligated into a NotI cut pWE15 cosmid, resulting in p $\beta$ LG-hLFgen (Fig. 19). The 34 kb NotI insert was isolated from the cosmid by NotI  
20 digestion and microinjected following standard procedures.

For the generation of p $\beta$ LG-hLFgen37 the 9.4 kb NotI-MluI fragment from p $\beta$ LGhLFgen9k was ligated with the 30 kb 3' MluI-ClaI fragment from hLF clone 13.1, combined with  
25 a ClaI-NotI linker into a NotI cut pWE15 cosmid vector.

The  $\beta$ LG-hLFgen insert was isolated and microinjected following standard procedure.

Expression data:

$\beta$ LG-hLFgen (the shorter of the 2 constructs) was  
30 injected and 7 independent mouse lines were produced. Expression data for hLF product in milk is available for the following lines:

-105-

<u>Construct</u>	<u>Line</u>	<u>Max. hLF expression in milk (mg/ml)</u>
$\beta$ LG-hLFgen	1106	0.02
	1107	1.9
	1108	0.8
	1110	6.2
	1111	1.3
	1155	2.1
	1156	2.2
	1591	0.05
$\beta$ LG-hLFgen37	1592	27
	1593	5.9

UBSTITUTE SHEET

-106-

EXAMPLE 21Isolation of a Genomic hLF Fragment  
Containing Both the Structural Gene  
and the hLF Promoter

5 HLF is normally expressed at relatively high levels (1-  
2 mg/ml) in human milk. ~~To determine whether the hLF~~  
~~promoter can drive high level hLF expression in the milk~~  
~~of transgenic animals, the intact hLF gene under control~~  
~~of its own promoter was microinjected using standard~~  
10 ~~procedures.~~

Construction details:

Two important points determined the construction route.  
Since the cosmid vector C2RB (Fig. 13) containing the  
genomic hLF clones does not contain unique restriction  
15 sites flanking the hLF insert, the intact insert could  
not be isolated directly from this cosmid. It was  
desirable to include all 5' and 3' flanking sequences  
present in hLF clones 13.1 and 13.2 into the transgene.  
Since clone 13.2 (Fig. 13) contains the most 5' flanking  
20 sequences (13 kb) and clone 13.1 the most 3' flanking  
sequences (7 kb more than 13.2), the 5' part of 13.2 was  
combined with the 3' part of 13.1.

The cosmid 13.2 was linearized at the PvuI site 0.5 to  
0.8 kb upstream of the 5' region of the hLF insert (Fig.  
25 20) and subsequently treated with the exonuclease Bal31,  
thereby removing approximately 1 kb of cosmid and 0.2  
to 0.5 of 5' hLF sequence. Subsequently, the DNA was  
treated with T4 polymerase to create blunt ends and cut  
with MluI. The approximately 19 kb (12.5 5' flanking  
30 sequences + 6.2 kb hLF gene) blunt end-MluI cut plasmid  
vector (pKUN6deltaCla, SmaI-MluI), resulted in plasmid  
phLF5'M gene 37. This plasmid contains a NotI site  
directly 5' of the SmaI site. From this plasmid, the  
19 kb NotI-MluI fragment was isolated and ligated with

-107-

the 30 kb MluI-NotI 3' fragment from construct 8hLFgen37 into a NotI cut pWE15 cosmid, resulting in p5'hLFgen37 (Fig. 20).

The 49 kb NotI insert was isolated and microinjected following standard procedures.

Expression data on construct p5'hLFgen37:

Eight independent founder mice have been generated for the p5'hLFgen37 construct; expression data are available for 6 lines.

10

Construct	Line	Max. hLF expression in milk (mg/ml)
<u>p5'hLFgen37</u>	1491	ND
	1492	2.5
	1493	4.2
	1495	6.5
	1496	18
	1497	ND
	1506	6.3
	1551	6.4

ND = not done

15

EXAMPLE 22

Generation of Mammary Gland  
Specific hLZ Expression Cassettes

The structure and sequence of the human lysozyme gene has been described (Peters, et al. (1989) Eur. J. Biochem  
20 182:507-516). The structural hLZ gene contains 4 exons and is 5.3 kb in size.

Using a 91-mer synthetic DNA sequence complementary to part of exon 2 of the hLZ gene as a probe, several

**SUBSTITUTE SHEET**

-108-

independent hLZ clones were isolated from a human genomic phage library. The clone  $\lambda$ 7.2.1 contains 14 kb insert comprising 8.7 kb of 5' flanking sequences and 5.3 kb of the genomic hLZ gene. Exon 4 is only partly included: clone  $\lambda$ 7.2.1 stops at one of the Sau3A sites at position 5333 and 5350 (numbering according to Peters, et al., op. cit.). The region downstream of position 5333/5350 (532 or 549 bp of exon 4 sequences) is missing. These sequences are non-coding and represent part of the 3' UTR of the hLZ gene. All hLZ coding sequences are present in  $\lambda$ 7.2.1.

#### Expression Vector 16,8hLZ

The design of expression vector 16,8hLZ, shown in Fig. 21, is as follows. The 5' flanking region (including the promoter) of the hLZ gene was removed and replaced with the bovine  $\alpha$ S1 casein gene promoter by subcloning into the plasmid p-16kbCS which is described in Example 16. The fusion site is located in the 5' UTR of the hLZ gene (exon 1), such that in addition to 23 bp of casein 5' UTR most of the hLZ 5' UTR is present. All coding sequences in this construct, including the signal sequence, are derived from hLZ clone  $\lambda$ 7.2.1 (Fig. 23A).

The 3'UTR of the hLZ gene in clone  $\lambda$ 7.2.1 was fused to the 3'UTR + flanking region of the bovine  $\alpha$ S1 casein gene described previously. The resulting 3'UTR of construct 16,8hLZ is therefore derived partly from the hLZ gene (exon 4, running from bp 4761 to bp 5333/5350) and partly (including part of exon 8 and all of exon 9) from the bovine  $\alpha$ S1 casein gene. The 3' flanking region (8 kb) is derived entirely from the bovine  $\alpha$ S1 casein gene.

-109-

Construction details for 16,8hLZ:16,hLZ:

The 6 bp directly 5' to the AUG codon in hLZ exon 1 constitute a HincII site. A SalI phage polylinker site is located directly 3' of the  $\lambda$ 7.2.1 insert. These sites were used to isolate a 5.3 kb HincII-SalI insert (Fig. 23). The sequence running from +3 (relative to the transcription initiation site at +1) to the HincII site, was synthesized by annealing the 31-mer and 35-mer depicted in Fig. 23A. The resulting synthetic DNA fragment has artificial 5' KpnI-HincII fragment and the 5.3 kb HincII-SalI fragment were subcloned into a KpnI-SalI cut pKUN-1 plasmid (Fig. 23A). From the resulting 9.3 kb plasmid (pKHLys3'5.3) the 5.3 kb ClaI-SalI fragment was isolated and subcloned into a Cla-SalI cut p-0.7kbCS plasmid (the equivalent of p-16CS but containing less 5' flanking sequences), resulting in pKhLZ0.7.

The 8 kb bovine  $\alpha$ S1 casein gene EcoRI fragment containing the 3' casein UTR and ca 6.6 kb of flanking sequences, was isolated from plasmid pKE3'E10 (described previously) as an 8 kb 5'-XhoI-SalI-3' fragment (Fig. 23B). This fragment was subcloned into the SalI site of pKhLZ0.7, resulting in p0.7,8hLZ. After this, the SalI site of p0.7,8hLZ was replaced with a NotI site by insertion of linker S1/S2 (Fig. 23C), yielding plasmid p0.7,8hLZnt (Fig. 23D). From this plasmid, the 13.3 kb ClaI-NotI fragment was isolated and ligated with the 14.5 kb NotI-ClaI fragment from p-16CS into a NotI cut pWE15 cosmid (Fig. 23E). From the resulting construct (named 16,8hLZ in Fig. 23E) the 27.8 kb NotI insert was isolated, purified and microinjected into murine and bovine zygotes following standard procedures.

-110-

Expression Vector 16,8hLZ3

The design of expression vector 16,8hLZ3, shown in Fig. 22, is as follows. Previously described expression vector 16,8hLF3 was used in the construction of

5 16,8hLZ3. The vector 16,8hLZ3 contains not only the bovine  $\alpha$ S1 casein gene promoter, but also the complete first exon and part of the first intron of the bovine  $\alpha$ S1 gene. In addition, it contains part of the first intron plus the splice acceptor site of an

10 immunoglobulin gene. The signal sequence and part of the 3' UTR and the complete 3' flanking region are also derived from the bovine  $\alpha$ S1 casein gene. The hLF cDNA and the  $\alpha$ S1 casein signal sequence are excised from this vector by ClaI-SalI double digestion. The ClaI site is

15 located 5 bp 5' to the translation initiation codon.

An 5.3 kb ClaI-SalI hLZ fragment was isolated from plasmid pKhLZ0.7 and subcloned into a ClaI-SalI cut 16,8hLF3 vector from which the hLF cDNA had been removed by ClaI-SalI double digestion.

20 The 16,8hLZ expression cassette vector sequence was removed by NotI digestion, subsequently purified according to standard procedures and microinjected into mouse zygotes.

Expression data:25 Construct 16,8hLZ:

Seven transgenic mice were generated for construct 16,8hLZ. Expression data are available for 6 independent mouse lines (data from lactating offspring, using our standard hLZ assay on milk samples).



-111-

<u>Construct</u>	<u>Line</u>	<u>Max. hLF expression in milk (mg/ml)</u>
16,8 hLZ	645	10
	647	0.7
	661	260
	662	7.4
	1069	60
	1070	28

The above data illustrates that 16,8hLZ expresses at relatively high levels. In human milk, hLZ levels are only 50 µg/ml (max). Since hLZ is a 15 kD protein, a level of 0.26 mg/ml hLZ compares to ~ 1.3 mg/ml of hLF (hLF is 80 kD).

Construct 16,8hLZ3:

Four independent transgenic mice were generated for covalent 16,8hLZ3. The following expression data are available from mouse lines 905 and 907.

<u>Mouse line</u>	<u>Expression (µg/ml) (max)</u>
905	475
907	10

The data show that 16,8hLZ3 can be expressed at relatively high levels (0.36 mg/ml compares to ~ 1.8 mg/ml hLF). However, as also shown, 16,8hLZ3 does not always express at high levels. Although the number of mice analyzed is very low, constructs 16,8hLZ and 16,8hLZ3 seem to behave more or less similar with regard to frequency of expression and expression levels. It should be noted, however, that another 7 lines of mice transgenic for 16,8hLZ also contain the 16,8hLF3 construct. (See below.) None of these lines expressed

**UBSTITUTE SHEET**

-112-

as high as 0.36 mg/ml. Therefore, 16,8hLZ3 appears to be a more efficient construct than 16,8hLZ. This could be caused by the heterologous splice site (which does enhance hLF cDNA expression levels).

5

EXAMPLE 23

Transgenic Mice Containing Transgenes  
Encoding Genomic hLZ and hLF cDNA

Coinjection of 16,8hLF3 and 16,8hLZ

To assess the feasibility of simultaneously expressing  
 10 hLF and hLZ in the milk of transgenic animals, the  
appropriate isolated and purified 16,8hLF3 and 16,8hLZ  
constructs were coinjected into murine zygotes.

Seven independent mouse lines transgenic for both  
 15 constructs were generated. The expression data  
 available for each line are as follows:

	<u>Mouse</u> <u>line</u>	<u>hLZ expression</u> <u>(<math>\mu</math>g/ml)</u>	<u>hLF expression</u> <u>(<math>\mu</math>g/ml)</u>
	649	150-250 (max: 311)	500-2000 (max: 2100)
20	650	10-30	1-9
	651	1-2.5	1-4.3
	657	1-6	1-15
	658	0.5	1
	659	<0.1	0.1
25	660	5-25	300-1260

Conclusions:

Only line 649 (1/7) expresses hLZ at relatively high levels. Line 649 and mouse line 660 (2/7) express high levels of hLF.

30 Comparison to data obtained  
from single construct injections:

For 16,8hLZ:

-113-

The hLZ expression level of mouse line 649, coinjected with the 16,8hLF3 and 16,8hLZ expression cassettes is comparable to that of line 661 injected only with 16,8hLZ.

5 In most cases, high level expression of hLZ is not obtained upon coinjection (1/7: high expression (line 649); 2/7 (650 and 660): intermediate-low expression; 4/7: low expression). Upon injection of single hLZ transgene, similar data are obtained (1/4: high  
10 expression (line 661); 1/4: intermediate; 2/4: low expression). Therefore, behavior of the 16,8hLZ transgene is not measurably influenced by the presence of the 16,8hLF3 transgene.

Note that none of the 7 lines expressed as high as line  
15 905 (construct 16,8hLZ3), although the level of 649 is in the same range.

In conclusion, these constructs can be expressed at relatively high levels (0.2-0.5 mg/ml range) with approximately 20-25 % of the resulting transgenic mice  
20 expressing at these high levels (3/13; 7 coinjections + 6 single inj.). Also coinjection with 16,8hLF3 does not appear to influence expression of 16,8hLZ.

For 16,8hLF3:

The single injection of 16,8hLF3 resulted in 13  
25 independent transgenic mouse lines which can be divided into 2 groups:

- (1) the low expressors which produced levels are from 0.1 to 5  $\mu$ g/ml (8/13) and
- (2) the high expressors which produced levels from  
30 40 to 200  $\mu$ g/ml (5/13).

Of the mice having the coinjected fragments, 2/7 express at high levels. This is similar to the frequency of

-114-

high level expression observed upon injection of one fragment (5/13). However, both 16,8hLF3/16,8hLZ mice lines (649 and 660) express hLF at much higher levels than observed previously. This indicates that the presence of the hLZ construct stimulates expression of the 16,8hLF3 construct. In line 649, the high hLF levels are accompanied by high hLZ levels. For line 660, this is less clear as hLZ levels are intermediate. However, as illustrated below, RNA analysis reveals that the 16,8hLZ transgene in line 660 is transcriptionally at least as active as the hLF transgene.

Results from expression analysis at the mRNA level:

Northern blot analysis was performed on both bovine lactating mammary gland total RNA and mammary gland total RNA from lactating transgenic mice (including mice transgenic for genomic hLF, 16,8hLF3 and 16,8hLF3 + 16,8hLZ). A 24 bp synthetic oligomer which hybridizes to exactly the same sequence in the 5' UTR of bovine  $\alpha$ S1 casein RNA and in all transgene derived RNA was used as a probe. Expression levels were compared directly by quantification of the amount of labelled probe hybridized to the transgene- and bovine  $\alpha$ S1 RNA.

It appeared that the ratio of hLZ- to hLF-mRNA and of hLZ- to bovine  $\alpha$ S1 mRNA was much higher than expected from the hLZ- and hLF protein levels. For example, line 649 expressed ~0.2 mg/ml of hLZ and ~1-2 mg/ml of hLF. After correcting for protein size (factor 5), hLZ and hLF mRNA levels are expected to be within the same range, with hLF levels about 2-fold higher than hLZ RNA levels. However, in line 649 hLZ mRNA levels were 20-fold higher than the hLF mRNA levels. Comparative RNA analysis of lines 650, 661, 662 and bovine mammary gland RNA confirmed these data.

-115-

It can therefore be concluded that transcriptionally, very high levels of hLZ expression are obtained using genomic hLZ sequences and the bovine  $\alpha$ S1 casein gene based expression system of the invention. The genomic hLZ constructs are transcribed at much higher levels than the hLF cDNA constructs, and expressed in the same range as the genomic hLF transgenes.

To compare the performance of different hLF and hLZ transgenes at the translational level, a 20-fold correction should be made. The transcriptional activity of hLZ transgenes expressing at 0.25 mg/ml is comparable to a protein level of 5 mg/ml, a level of 50  $\mu$ g/ml equivalent to 1 mg/ml. In addition, mouse line 649 hLZ mRNA levels exceeded bovine  $\alpha$ S1 mRNA levels - which had been 10-fold diluted - several fold. Since bovine  $\alpha$ S - casein is expressed at ~ 12 mg/ml (and is of similar size as hLZ), these hLZ RNA levels would be equivalent to an expression level of several mg/ml.

## EXAMPLE 24

20 Generation of 16,8 A hLZ:

Construct 16,8 A hLZ3 is a derivative of 16,8 hLZ3. In 16.8 A hLZ3 the hLZ 5' UTR sequences and the hLZ signal sequence have been replaced with the corresponding sequences from the bovine  $\alpha$ S1-casein gene.

25 Construct 16,A hLZ3 is a derivative of 16,8 A hLZ3. In 16.A hLZ3 the bovine  $\alpha$ S1-casein gene 3' UTR and flanking sequences have been replaced with the hLZ3' UTR and 4.5 kb of hLZ3" flanking sequences.

## Construction details:

30 Vector p07.8hLZ (Fig. 23B) was digested with ClaI and Sall. The 4.7 kb fragment (comprising 0.7 kb of the  $\alpha$ S1-casein 5' flanking sequences and the plasmid vector) was isolated and ligated to linker GP 278/279 (Fig. 25). This DNA sequence comprises part of the

-116-

bovine  $\alpha$ S1-casein 5' UTR, the complete bovine  $\alpha$ S1-casein signal sequence and 25 bp of hLZ sequence, encoding the N-terminal region of adult hLZ. The ligation product was isolated and ligated to a 5.3 kb Ball-Sall fragment from pKHLys3'5.3 (which is depicted in Fig. 23A). The resulting constructs is p0.7AhLZ $\Delta$ 3'. From this construct the 5.3 kb ClaI-Sall fragment was isolated and inserted into a ClaI-Sall vector, derived from p16,8hLF3 (also used in construction of 16,8 hLZ3). The resulting construct is designated p16,8A hLZ3 (Fig. 26).

For construction of 16,A hLZ3 the vector p0.7AhLZ $\Delta$ 3' was digested with XbaI and Sall, and a XbaI-NotI-Sall linker was inserted (Fig. 27). This vector was linearized with XbaI, and the 6.5 kb XbaI fragment from  $\lambda$ HLYS1 (described by Peters et al.; Eur. J. Biochem. 182, 507-516, 1989) was inserted in the sense orientation. This resulted in vector p0.7AhLZ. From this vector, the 9.8 kb ClaI-NotI hLZ fragment was isolated and, together with the 14.5 kb NotI-ClaI fragment from p16.8hLZ3, inserted into the NotI digested pWE15 cosmid vector.

In both cases, the transgene without plasmid sequences was isolated as a NotI fragment (16,8A hLZ3; 27.8 kb; 16,AhLZ3; 24.3 kb), purified and microinjected into fertilized mouse oocytes following standard procedures.

Four independent transgenic founder mice have been generated with construct 16,8 A hLZ3 and 6 mice were generated with construct 16 A hLZ3. Preliminary expression data:

	Mouse line	max. hLZ expression ( $\mu$ g/ml in milk)
16,8 A hLZ3:	1711	56
	1783	20
16 A hLZ3:	1806	267
	1809	2400

-117-

From these results it can be concluded that construct 16 A hLZ3 yields much higher expression levels than any other hLZ construct tested.

5 Preliminary quantitative Northern blotting data combined with data on hLZ protein expression levels indicate that the discrepancy between RNA and protein levels as observed for constructs 16,8hLZ and 16,8hLZ3 does not occur with construct 16 A hLZ3.

## EXAMPLE 25

10 Transmission experiments with "Calf #4"

Three heifers were super-ovulated using normal procedures used in cattle breeding (described in Diekman, S.J. et al. (1989) Theriogenology 31:473-487). These animals were subsequently inseminated with sperm  
15 from "calf #4" as described in Example 15. Calf #4 was judged to be transgenic as described in Example 15. The insemination resulted in two pregnancies.

These two animals were slaughtered four weeks after insemination and the embryos recovered from the uterus.  
20 Total DNA was isolated from these embryos following procedures as described in Maniatis et al. (1982), digested with EcoRI, and analyzed by "Southern Blot technique". The blot was hybridized to a probe specific for the hLF gene (same protocol as in example 15). Of  
25 the 12 embryos recovered 6 (50%) showed an hLF-specific band. In all cases the band was of the expected size and intensity. This indicates that:

- (a) the transgene transmits with an efficiency of appr. 50%
- 30 (b) the copy number is the same as in the founder (~3)

-118-

(c) no gross rearrangements have occurred during transmission

Of the six transgenic embryos, five were male and one was female according to a PCR-analysis with primers specific for a bovine Y-chromosome repeat. These data demonstrate that the transgene can be transmitted to both males and females and has not integrated in the Y-chromosome.

The sequences of the Y-chromosome specific primers are:

Forward primer: 5'-GGA TCC GAG ACA CAG AAC AGG-3'

Reverse primer: 5'-GCT AAT CCA TCC ATC CTA TAG-3'

#### EXAMPLE 26

##### Expression of recombinant proteins in saliva of calves

Ten animals were born from oocytes co-injected with the hLF transgene (as described in Example 18) and a hLZ transgene (16,8hLZ; described in Example 22). None of these animals appeared transgenic as judged by Southern Blot, but four of them (all males) were judged to be mosaic based on PCR with 0.5 µg DNA from blood and ear. Primers for this PCR-experiment were located in exon 8 of the hLF gene. The sequences of the primers are 5'-TTT GGA AAG GAC AAG TCA CCG-3' and 5'-CTC ACT TTT CCT CAA GTT CTG-3'

All ten animals were tested for hLF and hLZ expression in saliva. Epithelial cells in the salivary gland are structurally and functionally similar to such cells in the mammary gland, and some milk protein genes may also be expressed in salivary gland (albeit at much lower levels than in mammary gland).



-119-

Approximately 2 ml of saliva was collected from the mouth of the animal and levels of protein were determined in these samples using a radioimmunoassay as described in Example 5. Of the ten animals, three showed expression of hLF above the lower limit of detection. All three animals were part of the group of four animals judged to be mosaic.

Expression levels were as follows:

	animal	sample 1 (ng/ml)	sample 2 (ng/ml)
10	9772	25	18
	9773	3	1.4
	9774	1.2	nd

nd = not determined

All 10 animals were also tested for hLZ expression. Only animal 9772 showed expression of hLZ in saliva. The amount detected was 2 ng/ml.

Of the 21 animals born in the experiment described in Example 15, one animal (male) was judged to be mosaic based on the fact that it was immunotolerant for hLF. This animal showed an hLF expression in saliva of 100 ng/ml.

These data show that the transgenes used are capable of expressing hLF (and hLZ) in bovines.

Having described the preferred embodiments of the present invention, it will appear to those ordinarily skilled in the art that various modifications may be made to the disclosed embodiments, and that such modifications are intended to be within the scope of the present invention.

All references disclosed herein are expressly incorporated by reference.

-120-

WHAT IS CLAIMED IS:

1. A transgene for producing a recombinant polypeptide in transgenic bovine species comprising at least one expression regulation DNA sequence functional in at least one cell-type of said bovine species operably  
5 linked to a recombinant DNA encoding a recombinant polypeptide, wherein said transgene is capable of directing the expression of said recombinant DNA sequence in at least said one cell-type of a bovine species containing said transgene to produce said  
10 recombinant polypeptide.
2. The transgene of Claim 1 wherein said expression regulation sequences comprise 5' and 3' expression regulation sequences from a serum albumin, said cell-type is liver cell, said recombinant polypeptide is  
15 human serum albumin and said transgene further comprises a secretory DNA sequence functional in said liver cells and operably linked to the recombinant DNA encoding said human serum albumin.
3. A transgene for producing a recombinant polypeptide  
20 in the milk of transgenic bovine species comprising at least one expression regulation DNA sequence functional in the mammary secretory cells of said bovine species, a secretory DNA sequence encoding a secretory signal sequence also functional in the mammary secretory cells  
25 of said bovine species and a recombinant DNA sequence encoding a recombinant polypeptide, wherein said secretory DNA sequence is operably linked to said recombinant DNA sequence and to form a secretory-recombinant DNA sequence said at least one expression  
30 regulation sequence operably linked to said secretory-recombinant DNA sequence, such that said transgene is capable of directing the expression of said secretory-recombinant DNA sequence in mammary secretory cells of

-121-

bovine species containing said transgene to produce a form of recombinant polypeptide which when secreted from said mammary secretory cells produces recombinant polypeptide in the milk of said bovine species.

- 5           4.    The transgene of Claim 1 or 3 further comprising a recombinant intervening sequence.
5.    The transgene of Claim 4 wherein said recombinant intervening sequence is a hybrid intervening sequence.
- 10          6.    The transgene of Claim 5 wherein said hybrid intervening sequence contains a permissive RNA splice signal.
7.    The transgene of Claim 3 wherein said recombinant polypeptide is a homologous polypeptide from bovine species.
- 15          8.    The transgene of Claim 7 wherein said homologous polypeptide is selected from the group consisting of caseins, lactoferrin, lysozyme, cholesterol hydrolase and serum albumin.
- 20          9.    The transgene of Claim 3 wherein said recombinant polypeptide is a heterologous polypeptide.
10.   The transgene of Claim 9 wherein said heterologous polypeptide is selected from the group consisting of human milk proteins, human serum proteins, and industrial enzymes.
- 25          11.   The transgene of Claim 10 wherein said heterologous polypeptide is a human milk protein.
12.   The transgene of Claim 11 wherein said human milk protein is selected from the group consisting of

-122-

secretory immunoglobulins, lysozyme, lactoferrin, lactoglobulin,  $\alpha$ -lactalbumin and bile salt-stimulated lipase.

5 13. The transgene of Claim 12 wherein said milk protein is lactoferrin or lysozyme.

14. The transgene of Claim 10 wherein said heterologous polypeptide is a human serum protein.

10 15. The transgene of Claim 14 wherein said human serum protein is selected from the group consisting of albumin, immunoglobulin, Factor VIII, Factor IX and Protein C.

16. The transgene of Claim 15 wherein said serum protein is albumin.

15 17. The transgene of Claim 10 wherein said heterologous polypeptide is an industrial enzyme selected from the group consisting of proteases, lipases, chitinases and ligninases.

20 18. The transgene of Claim 3 wherein said secretory DNA sequence is selected from the group consisting of DNA sequences encoding secretory signal sequences from human lactoferrin, human serum albumin, human lysozyme and secretory signal sequences from bovine  $\alpha$ S1-casein,  $\alpha$ S2-casein,  $\beta$ -casein,  $\kappa$ -casein,  $\alpha$ -lactalbumin,  $\beta$ -lactoglobulin, and serum albumin.

25 19. The transgene of Claim 18 wherein said secretory DNA sequence is the DNA sequence encoding the signal secretion sequence of bovine  $\alpha$ S1 casein.

20. The transgene of Claim 3 wherein said at least one expression regulation sequence comprises 5' expression

-123-

regulation DNA sequences operably linked to the 5' end of said secretory-recombinant DNA sequence.

5 21. The transgene of Claim 20 wherein said 5' expression regulation DNA sequence is selected from the group consisting of 5' expression regulation sequence from bovine genes encoding  $\alpha$ S1-casein,  $\alpha$ S2-casein,  $\beta$ -casein,  $\kappa$ -casein,  $\alpha$ -lactalbumin, and  $\beta$ -lactoglobulin.

10 22. The transgene of Claim 21 wherein said 5' expression regulation DNA sequence is a proximal 5' expression regulation sequence comprising the promoter of bovine  $\alpha$ S1-casein.

15 23. The transgene of Claim 22 wherein said 5' expression regulation DNA sequence further comprises a distal 5' expression regulation sequence comprising 5'-flanking DNA sequence from bovine  $\alpha$ S1-casein.

24. The transgene of Claim 20 further comprising 3' expression regulation sequences operably linked to the 3' end of said secretory-recombinant DNA sequence.

20 25. The transgene of Claim 24 wherein said 3' expression regulation sequence comprise 3' expression regulation sequence from bovine genes encoding  $\alpha$ S1-casein,  $\alpha$ S2-casein,  $\beta$ -casein,  $\kappa$ -casein,  $\alpha$ -lactalbumin, and  $\beta$ -lactoglobulin.

25 26. The transgene of Claim 25 wherein said 3' expression regulation DNA sequence comprises a 3' proximal expression regulation sequence from bovine  $\alpha$ S1-casein.

27. The transgene of Claim 26 wherein said 3' expression regulation DNA sequence further comprises a

-124-

3' distal expression regulation sequence from bovine  
 $\alpha$ S1-casein.

5 28. The transgene of Claim 27 wherein said distal 5'  
expression regulation DNA sequence comprises about a  
30 kb 5'-flanking region of bovine  $\alpha$ S1-casein and said  
distal 3' expression regulation DNA sequence comprises  
about a 15 kb 3'-flanking region of bovine  $\alpha$ S1-casein.

10 29. A transgenic bovine species capable of producing  
a recombinant polypeptide in at least one cell type of  
said animal.

30. A transgenic bovine species capable of producing  
recombinant polypeptide in the milk of said transgenic  
species.

15 31. The transgenic bovine species of Claim 30 wherein  
said recombinant polypeptide is a homologous polypeptide  
from bovine species.

32. The transgenic bovine species of Claim 30 wherein  
said recombinant polypeptide is a heterologous  
polypeptide.

20 33. The transgenic bovine species of Claim 32 wherein  
said heterologous polypeptide is selected from the group  
consisting of human milk proteins, human serum proteins,  
and industrial enzymes..

25 34. The transgenic bovine species of Claim 33 wherein  
said heterologous polypeptide is a human milk protein.

35. The transgenic bovine species of Claim 34 wherein  
said human milk protein is selected from the group  
consisting of secretory immunoglobulins, lysozyme,

-125-

lactoferrin, lactoglobulin,  $\alpha$ -lactalbumin and bile salt-stimulated lipase.

36. The transgenic bovine species of Claim 35 wherein said milk protein is lactoferrin or lysozyme.

5 37. The transgenic bovine species of Claim 33 wherein said heterologous polypeptide is a human serum protein.

38. The transgenic bovine species of Claim 37 wherein said human serum protein is selected from the group consisting of albumin, immunoglobulin, Factor VIII, Factor IX and Protein C.

10

39. The transgenic bovine species of Claim 38 wherein said serum protein is albumin.

40. The transgenic bovine species of Claim 33 wherein said heterologous polypeptide is an industrial enzyme selected from the group consisting of proteases, lipases, chitinases and ligninases.

15

41. Milk from transgenic bovine species containing a recombinant polypeptide.

42. The milk of Claim 41 wherein said recombinant polypeptide is a homologous polypeptide from bovine species.

20

43. The milk of Claim 41 wherein said recombinant polypeptide is a heterologous polypeptide.

44. The milk of Claim 43 wherein said heterologous polypeptide is selected from the group consisting of human milk proteins, human serum proteins, and industrial enzymes.

25

-126-

45. The milk of Claim 44 wherein said heterologous polypeptide is a human milk protein.

5        46. The milk of Claim 45 wherein said human milk protein is selected from the group consisting of secretory immunoglobulins, lysozyme, lactoferrin, lactoglobulin,  $\alpha$ -lactalbumin and bile salt-stimulated lipase.

47. The milk of Claim 46 wherein said milk protein is lactoferrin or lysozyme.

10       48. The milk of Claim 43 wherein said heterologous polypeptide is a human serum protein.

15       49. The milk of Claim 48 wherein said human serum protein is selected from the group consisting of albumin, immunoglobulin, Factor VIII, Factor IX and Protein C.

50. The milk of Claim 49 wherein said serum protein is albumin.

51. A food formulation comprising transgenic milk containing a recombinant polypeptide.

20       52. The food formulation of Claim 51 wherein said recombinant polypeptide is at least partially purified from said transgenic milk.

53. The food formulation of Claim 51 formulated with nutrients appropriate for infant formula.

25       54. A method for producing a transgenic bovine species capable of producing a recombinant polypeptide in the milk of said bovine species, said method comprising:



-127-

introducing the transgene of Claim 1 into an embryonal target cell of a bovine species;

transplanting the transgenic embryonal target cell formed thereby or the embryo obtained herefrom into a recipient female bovine parent; and

identifying at least one female offspring which is capable of producing said recombinant polypeptide in the milk of said offspring.

55. A method for producing a transgenic non-human mammal having a desirable phenotype comprising:

(a) methylating a transgene capable of conferring said phenotype when incorporated into the cells of said transgenic non-human animal;

(b) introducing said methylated transgene into fertilized oocytes of said non-human mammal to permit integration of said transgene into the genomic DNA of said fertilized oocytes;

(c) culturing the individual oocytes formed hereby to pre-implantation embryos thereby replicating the genome of each of said fertilized oocytes;

(d) removing at least one cell from each of said pre-implantation embryos and lysing said at least one cell to release the DNA contained therein;

(e) contacting said released DNA with a restriction endonuclease capable of cleaving said methylated transgene but incapable of cleaving the unmethylated form of said transgene formed after integration into and replication of said genomic DNA; and

(f) detecting which of said cells from said pre-implantation embryos contain a transgene which is resistant to cleavage by said restriction endonuclease as an indication of which pre-implantation embryos have integrated said transgene.

-128-

56. The method of Claim 55 wherein said removal of at least one cell forms a first and second hemi-embryo for each of said pre-implementation embryos and each of said first hemi-embryos is lysed and analyzed according to steps (d) through (f), said method further comprising;
- 5 (g) cloning at least one of said second hemi-embryos; and
- (h) to form a multiplicity of transgenic embryos.
- 10 57. The method of Claim 56 further comprising transplanting more than one of said transgenic embryos into recipient female parents to produce a population containing at least two transgenic non-human animals having the same genotype.
- 15 58. The method of Claim 55 further comprising transplanting the remainder of said pre-implantation embryo containing a genomically integrated transgene into a recipient female parent and identifying at least one offspring having said phenotype.
- 20 59. The method of Claim 55 wherein said restriction endonuclease is DPNI and said transgene is methylated at N6 of the adenine of the sequence GATC contained within said transgene.
- 25 60. The method of Claim 59 wherein said detection utilizes a polymerase chain reaction using extension primers complementary to sequences upstream and downstream to said GATC sequence.
- 30 61. The method of Claim 59 wherein said non-human transgenic mammal is bovine species, said transgene encodes a recombinant polypeptide and said desired phenotype is the ability to produce said recombinant polypeptide in the milk of said bovine species.

-129-

62. The method of Claim 61 wherein said transgene is the transgene of Claim 3.

63. A transgene for producing a recombinant polypeptide in the milk of transgenic bovine species comprising:

5 (i) a bovine 5' expression regulation sequence;

(ii) a secretory DNA sequence encoding a secretory signal sequence functional in the mammary secretory cells of the bovine species;

10 (iii) a recombinant DNA sequence encoding a recombinant polypeptide, said secretory DNA sequence being operably linked to said recombinant DNA sequence, wherein a secretory-recombinant DNA sequence is formed, said secretory-recombinant DNA sequence being operably  
15 linked to said bovine expression regulation sequence;

(iv) a 3' untranslated sequence;

(v) a 3' flanking sequence of a bovine gene;

and

20 wherein said transgene is capable of directing the expression of said secretory-recombinant DNA sequence in mammary secretory cells of bovine species containing said transgene

25 to produce a form of recombinant polypeptide which when secreted from said mammary secretory cells produces recombinant polypeptide in the milk of said bovine species.

64. The transgene of claim 63, further comprising a recombinant intervening sequence.

30 65. The transgene of claim 64 wherein the recombinant intervening sequence is a hybrid intervening sequence.

66. The transgene of claim 65 wherein the hybrid intervening sequence comprises a 5' portion of an intervening sequence from bovine  $\alpha$ -S<sub>1</sub>-casein and a 3'

-130-

sequence portion of an IgG heavy chain intervening sequence.

5 67. The transgene of claim 66 wherein the 3' sequence portion is a 3' splice signal sequence associated with the J-C segment rearrangement of an IgG heavy chain.

68. The transgene of claim 63, wherein the bovine expression regulation sequence and the 3' flanking sequence are derived from the same bovine gene.

10 69. The transgene of claim 63, wherein the bovine expression regulation sequence, the 3' untranslated sequence, and the 3' flanking sequence are derived from the same bovine gene.

70. The transgene of claim 68 or claim 69 wherein the bovine gene is  $\alpha$ -S<sub>1</sub>-casein.

15 71. The transgene of Claim 70 wherein the bovine expression regulation sequence comprises about a 30kb 5'-flanking region of bovine  $\alpha$ S1-casein and the 3'-flanking sequence comprises about a 15kb 3'-flanking region of bovine  $\alpha$ S1-casein.

20 72. The transgene of claim 3 or claim 63 wherein the milk comprises greater than 50 micrograms of the recombinant polypeptide per milliliter.

73. A transgenic bovine species capable of producing a recombinant polypeptide in saliva.

25 74. The semen of a transgenic bovine.

75. A transgene for producing a recombinant polypeptide in the milk of transgenic bovine species comprising:

-131-

(i) a 5' expression regulation sequence;  
(ii) a secretory DNA sequence encoding a secretory signal sequence functional in the mammary secretory cells of the bovine species;  
5 (iii) a recombinant DNA sequence encoding a recombinant polypeptide, said secretory DNA sequence being operably linked to said recombinant DNA sequence, wherein a secretory-recombinant DNA sequence is formed, said secretory-recombinant DNA sequence being operably  
10 linked to the 5' expression regulation sequence;  
(iv) a 3' untranslated sequence; and,  
(v) a 3' flanking sequence from a human gene;  
wherein the transgene is capable of directing the expression of the secretory-recombinant DNA sequence  
15 in mammary secretory cells of bovine species containing the transgene  
to produce a form of recombinant polypeptide which when secreted from the mammary secretory cells produces recombinant polypeptide in the milk of the bovine  
20 species.

76. The transgene of claim 75, wherein the 5' expression regulatory sequence is a bovine sequence.

77. The transgene of claim 75 or claim 76, wherein the 3' flanking sequence is from the human lactoferrin (hLF) gene.  
25

78. The transgene of claim 77, wherein the 3' flanking sequence is 9 kilobase pairs in length.

79. The transgene of claim 77, wherein the recombinant polypeptide is human lactoferrin.

80. The transgene of claim 75, wherein the 5' expression regulation sequence, the secretory DNA sequence, the recombinant DNA sequence encoding a  
30

-132-

recombinant polypeptide, the 3' untranslated sequence; and the 3' flanking sequence are from a human gene.

81. The transgene of claim 80, wherein the human gene is the lactoferrin gene.

5 82. A method for expressing a human polypeptide in the milk of a bovine comprising:

introducing a human genomic fragment encoding the human polypeptide into an embryonal target cell of a bovine species;

10 transplanting the transgenic embryonal target cell formed thereby or the embryo obtained therefrom into a recipient female bovine parent; and

15 identifying at least one female offspring which is capable of producing the recombinant polypeptide in the milk of the offspring.

83. The method of claim 82, wherein the human polypeptide is lactoferrin.

1/45

```

sauSEI
haeIII
asuI
sauSEI
nleIU
hgiIII
eccOIEI
baeIIEI
banII
asuI
nleIU
nnII
aval apal nnII
nnII eccOIEI
1 66A CTT CTC TTC CTC CTC CTC CTC TTC CTC 66G GCC CTC 66A CTC
-16 61y Leu Val Phe Leu Val Leu Leu Phe Leu 61y Ala Leu 61y Leu
haeIII
aaeI
cfrI
hinFI
hhaI
4E TBT CTC GCT GGC CGT AGG AGA AGG AGT GTT CAG TGG TGC GCC GTA TCC
-3 Cys Leu Ala 61y Arg Arg Arg Arg Ser Val 61n Trp Cys Ala Val Ser
haeIII
nnII
aval haeI
54 CAA CCC GAG GCC ACA AAA TCC TTC CAA TGG CAA AGG AAT ATG AGA AAA
14 61n Pro 61u Ala Thr Lys Cys Phe 61n Trp 61n Arg Asn Met Arg Lys
nnII
fnuSH-
sauSEI
baeI
pieI
haeIII
aaeI
hinFI
baeI
asuI
foiI
142 GTG CTT GGC CTT CTT GTC AGC TGC ATA AAG AGA GAC TCC CCC ATC CAG
30 Val Arg 61y Pro Pro Val Ser Cys 11e Lys Arg Asp Ser Pro 11e 61n
haeIII
haeI
scrFI
eccPI
batNI
haeIII
sauSEI
asuI
sfaNI
192 TBT ATC CAG GCC ATT GCG GAA AAC AGG GCC GAT GCT GTG ACC CTT GAT
4E Cys 11e 61n Ala 11e Ala 61u Asn Arg Ala Asp Ala Val Thr Leu Asp
sauSEI
nleIU
scrFI
eccRII
batNI
haeIII
stuI
haeIII
nnII
haeI
asuI
238 GGT GGT TTC ATA TAC GAG GCA GGC CTC GCC CCC TAC AAA CTC CGA CCT
82 61y 61y Phe 11e Tyr 61u Ala 61y Leu Ala Pro Tyr Lys Leu Arg Pro
sauSEI
avalI
asuI
fnuSHI
accI
nleIU
22E GTA GCG GCG GAA GTC TAC GGG ACC GAA AGA CAG CCA CGA ACT CACTAT
76 Val Ala Ala 61u Val Tyr 61y Thr 61u Arg 61n Pro Arg Thr His Tyr

```

Fig-1 Page 1 of 6

2/45

mbol1 {nu4H} alu1  
 hph1 bbv1  
 334 TAT GCG GTG GCT GTG GTG AAG AAG GGC GGC AGC TTT CAG CTG AAC GAA  
 94 Tyr Ala Val Ala Val Val Lys Lys Gly Gly Ser Phe Gln Leu Asn Glu

hae111 sau9E1  
 stu1 ave11  
 bpl1 hae1 esu1 fol1  
 362 CTG CAA GGT CTG AAG TCC TGC CAC ACA GGC CTT GCG AGG ACC GCT GGA  
 118 Leu Gln Gly Leu Lys Ser Cys His Thr Gly Leu Arg Arg Thr Ala Gly

sau9E1  
 ave11  
 esu1  
 nla1V  
 438 TGG AAT GTC CCT ACA GGG ACA CTT GGT CCA TTC TTG AAT TGG ACB GGT  
 126 Trp Asn Val Pro Thr Gly Thr Leu Arg Pro Phe Leu Asn Trp Thr Gly

hpl111 alu1  
 bap128E {nu4H}  
 ban11 bbv1 dde1 alu1  
 dde1 nhl1 pvu11 mbol1 pvu11  
 478 CCA CCT GAG CCC ATT GAG GCA GCT GTG CAG TTC TTC TCA GCG AGC TGT  
 142 Pro Pro Glu Pro Ile Glu Ala Ala Val Gln Phe Phe Ser Ala Ser Cys

mad1  
 hpe11  
 scrF1  
 nci1  
 sau11  
 528 GTT CCC GGT GCA GAT AAA GGA CAG TTC CCC AAC CTG TGT GCG CTG TGT  
 156 Val Pro Gly Ala Asp Lys Gly Gln Phe Pro Asn Leu Cys Arg Leu Cys

nla1V  
 scrF1  
 eccR11  
 nhl1 bsiN1 paa1  
 574 GCG GGG ACA GGG GAA AAC AAA TGT GCG TTC TCC TCC CAG GAA CCG TAC  
 174 Ala Gly Thr Gly Glu Asn Lys Cys Ala Phe Ser Ser Gln Glu Pro Tyr

nla1V  
 hpl11  
 ba-1 dde1 bana1 bana1  
 622 TTC AGC TAC TCT GGT GCG TTC AAG TGT CTG AGA GAC GGG GCT GGA GAC  
 182 Phe Ser Tyr Ser Gly Ala Phe Lys Cys Leu Arg Asp Gly Ala Gly Asp

sau9E1  
 ave11  
 esu1  
 bduM1  
 eccD1891  
 hpl111  
 bap128E nhl1 nhl1  
 678 GTG GCT TTT ATC AGA GAG AGC ACA GTG TTT GAG GAC CTG TCA GAC GAG  
 206 Val Ala Phe Ile Arg Glu Ser Thr Val Phe Glu Asp Leu Ser Asp Glu

718 GCT GAA AGG GAC GAG TAT GAG TTA CTC TGC CCA GAC AAC ACT GGG AAG  
 222 Ala Glu Arg Asp Glu Tyr Glu Leu Leu Cys Pro Asp Asn Thr Arg Lys

Fig-1 page 2 of 6



nseI  
 hpaII 3/45  
 cauI  
 nseI sauSEI  
 nseI nlaIV  
 scrFI  
 nciI eaeII  
 cauI  
 eaeI esuI  
 sauSEI ppmI  
 haeIII nlaIV  
 esuI ecoDI2E: nlaIII  
 7EE CCA GTG GAC AAG TTC AAA GAC TGC CAT CTG GCC CGG GTC CCT TCT CAT  
 23E Pro Val Asp Lys Phe Lys Asp Cys His Leu Ala Arg Val Pro Ser His  
 efaNI  
 folI mbdII  
 bglI draIII nmlI hinfI  
 814 GCC GTT GTG GCA CGA AGT GTG AAT GGC AAG GAG GAT GCC ATC TGG AAT  
 254 Ala Val Val Ala Arg Ser Val Asn Gly Lys Glu Asp Ala Ile Trp Asn  
 scrFI  
 scoRII  
 bstNI hphI  
 6E2 CTT CTC CGC CAG GCA CAG GAA AAG TTT GGA AAG GAC AAG TCA CCG AAA  
 27E Leu Leu Arg Gln Ala Gln Glu Lys Phe Gly Lys Asp Lys Ser Pro Lys  
 sau3AI  
 nciI  
 bsrI  
 ahoII  
 bstVI  
 bclII  
 510 TTC CAG CTC TTT GGC TCC CCT AGT GGG CAG AAA GAT CTG CTG TTC AAG  
 26E Phe Gln Leu Phe Gly Ser Pro Ser Gly Gln Lys Asp Leu Leu Phe Lys  
 nlaII  
 hpiCI  
 pjaI nmlI bspI2EE nmlI  
 hinfI taclI bsrI eaeI hinfI  
 5EE GAC TCT GGC ATT GGG TTT TCG AGG GTG CCC CGC AAG ATA GAT TCT GGG  
 32E Asp Ser Ala Ile Gly Phe Ser Arg Val Pro Pro Arg Ile Asp Ser Gly  
 nseI  
 styI hpaII  
 rsaI nlaIV folI nmlI  
 102E CTG TAC CTT GGC TCC GGC TAC TTC ACT GCC ATC CAG AAC TTG AAG AAA  
 31E Leu Tyr Leu Gly Ser Gly Tyr Phe Thr Ala Ile Gln Asn Leu Arg Lys  
 nspI  
 hpaII thaI  
 scrFI fnuDII  
 nciI bstUI  
 nmlI fnu4HI hinfI  
 nmlI bbvI cauII hhaI  
 1254 AGT GAG GAG GAA GTG GCT GCC CGG CGT GCG CGG GTC GTG TGG TGT GCG  
 334 Ser Glu Glu Glu Val Ala Ala Arg Arg Ala Arg Val Val Trp Cys Ala  
 hinfI  
 nmlI  
 fspI  
 fnu4HI  
 aluI hhaI bstXI  
 aluNI bbvI bsrI  
 1182 GTG GGC GAG CAG GAG CTG CGC AAG TGT AAC CAG TGG AGT GGC TTG AGC  
 35E Val Gly Glu Gln Glu Leu Arg Lys Cys Asn Gln Trp Ser Gly Leu Ser  
 fnu4HI nmlI  
 bbvI bspMI nmlI haeIII nmlI efaNI  
 115E GAA GGC AGC GTG ACC TGC TCC TCG GCC TCC ACC ACA GAG GAC TGC ATC  
 35E Glu Gly Ser Val Thr Cys Ser Ser Ala Ser Thr Thr Glu Asp Cys Ile

Fig 1 Page 3 of 6

4/45

```

scrFI
ecorII
batNI
          aluI  bfaNI nlaIII  folI  mnII
115E ECC CTG CTG CTG AAA GGA GAA GCT GAT GGC ATG AGT TTG GAT GGA GGA
38C Ala Leu Val Leu Lys Gly Glu Ala Asp Ala Met Ser Leu Asp Gly Gly

          nlaIII  nlaIV  scrFI
          bphI  hpcII  ecorII
          paaI  nspCII  banI  batNI
124E TAT GTG TAC ACT GCA TGC AAA TGT GGT TTG GTG CCT GTC CTG GCA GAG
35E Tyr Val Tyr Thr Ala Cys Lys Cys Gly Leu Val Pro Val Leu Ala Glu

          sau3AI
          mbcI
          dphI
          aluI
129A AAC TAC AAA TCC CAA CAA AGC AGT GAC CCT GAT CCT AAC TGT GTG GAT
41A Asn Tyr Lys Ser Gln Gln Ser Ser Asp Pro Asp Pro Asn Cys Val Asp

          sau3AI
          mbcI
          dphI
          ecorNI  ecorV
134C AGA CCT GTG GAA GGA TAT CTT GCT GTG GCG GTG GTT AGG AGA TCA GAC
43C Arg Pro Val Glu Gly Tyr Leu Ala Val Ala Val Val Arg Arg Ser Asp

scrFI
ecorII
batNI
129C ACT AGC CTT ACC TGG AAC TCT GTG AAA GGC AAG AAG TCC TGC CAC ACC
44E Thr Ser Leu Thr Thr Asn Ser Val Lys Gly Lys Lys Ser Cys His Thr

          haeIII
          nlaIII
          styI sau3EI  mbcII
          nccI asuI  earI
142E GGC GTG GAC AGG ACT GCA GGC TGG AAT ATC CCC ATG GGC CTG CTC TTC
45C Ala Val Asp Arg Thr Ala Gly Thr Asn Ile Pro Met Gly Leu Leu Phe

          nlaIV
          hpcIII
          bspICEE
          banII
          bspI
          aluI bspICEE
148E AAC CAG AGC GGC TCC TGC AAA TTT GAT GAA TAT TTC AGT CAA AGC TGT
47E Asn Gln Thr Gly Ser Cys Lys Phe Asp Glu Tyr Phe Ser Gln Ser Cys

          sau3AI
          mbcI
          dphI
          xhoII
          bspI
          hpaI
          bspI286
153A GGC CCT GGG TCT GAC CCG AGA TCT AAT CTC TGT GCT CTG TGT ATT GGC
48A Ala Pro Gly Ser Asp Pro Arg Ser Asn Leu Cys Ala Leu Cys Ile Gly

          hphI
          bspI286
158Z GAC GAG CAG GGT GAG AAT AAG TGC GTG CCC AAC AGC AAT GAG AGA TAC
51B Asp Glu Gln Gly Glu Asn Lys Cys Val Pro Asn Ser Asn Glu Arg Tyr

          nlaIV
          hpcI
          banI scrFI
          mspI  ecorII
          bspI  hpaII  batNI  ddeI  banI  bnaI
163C TAC GGC TAC ACT GGG GCT TTC CCG TGC CTG GCT GAG AAT GCT GGA GAC
52E Tyr Gly Tyr Thr Gly Ala Phe Arg Cys Leu Ala Glu Asn Ala Gly Asp

167B GTT GCA TTT GTG AAA GAT GTC ACT GTC TTG CAG AAC ACT GAT GGA AAT
54Z Val Ala Phe Val Lys Asp Val Thr Val Leu Gln Asn Thr Asp Gly Asn

```

Fig-1 Page 4 of 6



6/45

```

                                scrF1
                                ecor11
                                ba1h1
                                ng1A1
                                bsc12EE   m111   m111
2052 652 ATT ACT AAT CTG AAA AAG TGC TCA ACC TCC CCC CTC CTG GAA GGC
672 612 Jle Thr Asn Leu Lys Lys Cys Ser Thr Ser Pro Leu Leu Glu Ala

                                mde1
                                nat11
                                m111
                                ecor11
                                mbo11   sau3E1
                                mbo11   hae111
2112 TGT GAA TTC CTC AGG AAG TAA   mbo11   sau1 alu1
688 Cys Glu Phe Leu Arg Lys DC-   AACC6AAGAA GATG6GCCAG CTCCCCAAGA

                                b1y1
                                hae111
                                sau3E1
                                sau1
                                ecor11
                                m1e1V
                                ecor12E1 m1e1V
2181 AAGGCTCAGC CATTCACTGC CCCCAGTCTT TCTCCCCAGG TGTGTTG6GG CCTTGGCTCC
                                ecor1
                                fol1
                                mde1
2221 CCGTGTGAAA GTCGGGATTC CCGATCCATC TCTTACAA TCCCTGCTGT CTTCTAGCA
2281 AGAAGTAAAA TGAGAAATTT TTTTATATT CAAAAAAA

Length: 2218

```


Fig-1 page 6 of 6

7/45

1 3 5 7 9 11 13 15 17 19 21 23 25 27 29 31 33 35 37 39 41 43 45 47 49 51 53 55 57 59 61 63 65 67 69 71 73 75 77 79 81 83 85 87 89 91 93 95 97 99 101 103 105 107 109 111 113 115 117 119 121 123 125 127 129 131 133 135 137 139 141 143 145 147 149 151 153 155 157 159 161 163 165 167 169 171 173 175 177 179 181 183 185 187 189 191 193 195 197 199 201 203 205 207 209 211 213 215 217 219 221 223 225 227 229 231 233 235 237 239 241 243 245 247 249 251 253 255 257 259 261 263 265 267 269 271 273 275 277 279 281 283 285 287 289 291 293 295 297 299 301 303 305 307 309 311 313 315 317 319 321 323 325 327 329 331 333 335 337 339 341 343 345 347 349 351 353 355 357 359 361 363 365 367 369 371 373 375 377 379 381 383 385 387 389 391 393 395 397 399 401 403 405 407 409 411 413 415 417 419 421 423 425 427 429 431 433 435 437 439 441 443 445 447 449 451 453 455 457 459 461 463 465 467 469 471 473 475 477 479 481 483 485 487 489 491 493 495 497 499 501 503 505 507 509 511 513 515 517 519 521 523 525 527 529 531 533 535 537 539 541 543 545 547 549 551 553 555 557 559 561 563 565 567 569 571 573 575 577 579 581 583 585 587 589 591 593 595 597 599 601 603 605 607 609 611 613 615 617 619 621 623 625 627 629 631 633 635 637 639 641 643 645 647 649 651 653 655 657 659 661 663 665 667 669 671 673 675 677 679 681 683 685 687 689 691 693 695 697 699 701 703 705 707 709 711 713 715 717 719 721 723 725 727 729 731 733 735 737 739 741 743 745 747 749 751 753 755 757 759 761 763 765 767 769 771 773 775 777 779 781 783 785 787 789 791 793 795 797 799 801 803 805 807 809 811 813 815 817 819 821 823 825 827 829 831 833 835 837 839 841 843 845 847 849 851 853 855 857 859 861 863 865 867 869 871 873 875 877 879 881 883 885 887 889 891 893 895 897 899 901 903 905 907 909 911 913 915 917 919 921 923 925 927 929 931 933 935 937 939 941 943 945 947 949 951 953 955 957 959 961 963 965 967 969 971 973 975 977 979 981 983 985 987 989 991 993 995 997 999 1001 1003 1005 1007 1009 1011 1013 1015 1017 1019 1021 1023 1025 1027 1029 1031 1033 1035 1037 1039 1041 1043 1045 1047 1049 1051 1053 1055 1057 1059 1061 1063 1065 1067 1069 1071 1073 1075 1077 1079 1081 1083 1085 1087 1089 1091 1093 1095 1097 1099 1101 1103 1105 1107 1109 1111 1113 1115 1117 1119 1121 1123 1125 1127 1129 1131 1133 1135 1137 1139 1141 1143 1145 1147 1149 1151 1153 1155 1157 1159 1161 1163 1165 1167 1169 1171 1173 1175 1177 1179 1181 1183 1185 1187 1189 1191 1193 1195 1197 1199 1201 1203 1205 1207 1209 1211 1213 1215 1217 1219 1221 1223 1225 1227 1229 1231 1233 1235 1237 1239 1241 1243 1245 1247 1249 1251 1253 1255 1257 1259 1261 1263 1265 1267 1269 1271 1273 1275 1277 1279 1281 1283 1285 1287 1289 1291 1293 1295 1297 1299 1301 1303 1305 1307 1309 1311 1313 1315 1317 1319 1321 1323 1325 1327 1329 1331 1333 1335 1337 1339 1341 1343 1345 1347 1349 1351 1353 1355 1357 1359 1361 1363 1365 1367 1369 1371 1373 1375 1377 1379 1381 1383 1385 1387 1389 1391 1393 1395 1397 1399 1401 1403 1405 1407 1409 1411 1413 1415 1417 1419 1421 1423 1425 1427 1429 1431 1433 1435 1437 1439 1441 1443 1445 1447 1449 1451 1453 1455 1457 1459 1461 1463 1465 1467 1469 1471 1473 1475 1477 1479 1481 1483 1485 1487 1489 1491 1493 1495 1497 1499 1501 1503 1505 1507 1509 1511 1513 1515 1517 1519 1521 1523 1525 1527 1529 1531 1533 1535 1537 1539 1541 1543 1545 1547 1549 1551 1553 1555 1557 1559 1561 1563 1565 1567 1569 1571 1573 1575 1577 1579 1581 1583 1585 1587 1589 1591 1593 1595 1597 1599 1601 1603 1605 1607 1609 1611 1613 1615 1617 1619 1621 1623 1625 1627 1629 1631 1633 1635 1637 1639 1641 1643 1645 1647 1649 1651 1653 1655 1657 1659 1661 1663 1665 1667 1669 1671 1673 1675 1677 1679 1681 1683 1685 1687 1689 1691 1693 1695 1697 1699 1701 1703 1705 1707 1709 1711 1713 1715 1717 1719 1721 1723 1725 1727 1729 1731 1733 1735 1737 1739 1741 1743 1745 1747 1749 1751 1753 1755 1757 1759 1761 1763 1765 1767 1769 1771 1773 1775 1777 1779 1781 1783 1785 1787 1789 1791 1793 1795 1797 1799 1801 1803 1805 1807 1809 1811 1813 1815 1817 1819 1821 1823 1825 1827 1829 1831 1833 1835 1837 1839 1841 1843 1845 1847 1849 1851 1853 1855 1857 1859 1861 1863 1865 1867 1869 1871 1873 1875 1877 1879 1881 1883 1885 1887 1889 1891 1893 1895 1897 1899 1901 1903 1905 1907 1909 1911 1913 1915 1917 1919 1921 1923 1925 1927 1929 1931 1933 1935 1937 1939 1941 1943 1945 1947 1949 1951 1953 1955 1957 1959 1961 1963 1965 1967 1969 1971 1973 1975 1977 1979 1981 1983 1985 1987 1989 1991 1993 1995 1997 1999 2001 2003 2005 2007 2009 2011 2013 2015 2017 2019 2021 2023 2025 2027 2029 2031 2033 2035 2037 2039 2041 2043 2045 2047 2049 2051 2053 2055 2057 2059 2061 2063 2065 2067 2069 2071 2073 2075 2077 2079 2081 2083 2085 2087 2089 2091 2093 2095 2097 2099 2101 2103 2105 2107 2109 2111 2113 2115 2117 2119 2121 2123 2125 2127 2129 2131 2133 2135 2137 2139 2141 2143 2145 2147 2149 2151 2153 2155 2157 2159 2161 2163 2165 2167 2169 2171 2173 2175 2177 2179 2181 2183 2185 2187 2189 2191 2193 2195 2197 2199 2201 2203 2205 2207 2209 2211 2213 2215 2217 2219 2221 2223 2225 2227 2229 2231 2233 2235 2237 2239 2241 2243 2245 2247 2249 2251 2253 2255 2257 2259 2261 2263 2265 2267 2269 2271 2273 2275 2277 2279 2281 2283 2285 2287 2289 2291 2293 2295 2297 2299 2301 2303 2305 2307 2309 2311 2313 2315 2317 2319 2321 2323 2325 2327 2329 2331 2333 2335 2337 2339 2341 2343 2345 2347 2349 2351 2353 2355 2357 2359 2361 2363 2365 2367 2369 2371 2373 2375 2377 2379 2381 2383 2385 2387 2389 2391 2393 2395 2397 2399 2401 2403 2405 2407 2409 2411 2413 2415 2417 2419 2421 2423 2425 2427 2429 2431 2433 2435 2437 2439 2441 2443 2445 2447 2449 2451 2453 2455 2457 2459 2461 2463 2465 2467 2469 2471 2473 2475 2477 2479 2481 2483 2485 2487 2489 2491 2493 2495 2497 2499 2501 2503 2505 2507 2509 2511 2513 2515 2517 2519 2521 2523 2525 2527 2529 2531 2533 2535 2537 2539 2541 2543 2545 2547 2549 2551 2553 2555 2557 2559 2561 2563 2565 2567 2569 2571 2573 2575 2577 2579 2581 2583 2585 2587 2589 2591 2593 2595 2597 2599 2601 2603 2605 2607 2609 2611 2613 2615 2617 2619 2621 2623 2625 2627 2629 2631 2633 2635 2637 2639 2641 2643 2645 2647 2649 2651 2653 2655 2657 2659 2661 2663 2665 2667 2669 2671 2673 2675 2677 2679 2681 2683 2685 2687 2689 2691 2693 2695 2697 2699 2701 2703 2705 2707 2709 2711 2713 2715 2717 2719 2721 2723 2725 2727 2729 2731 2733 2735 2737 2739 2741 2743 2745 2747 2749 2751 2753 2755 2757 2759 2761 2763 2765 2767 2769 2771 2773 2775 2777 2779 2781 2783 2785 2787 2789 2791 2793 2795 2797 2799 2801 2803 2805 2807 2809 2811 2813 2815 2817 2819 2821 2823 2825 2827 2829 2831 2833 2835 2837 2839 2841 2843 2845 2847 2849 2851 2853 2855 2857 2859 2861 2863 2865 2867 2869 2871 2873 2875 2877 2879 2881 2883 2885 2887 2889 2891 2893 2895 2897 2899 2901 2903 2905 2907 2909 2911 2913 2915 2917 2919 2921 2923 2925 2927 2929 2931 2933 2935 2937 2939 2941 2943 2945 2947 2949 2951 2953 2955 2957 2959 2961 2963 2965 2967 2969 2971 2973 2975 2977 2979 2981 2983 2985 2987 2989 2991 2993 2995 2997 2999 3001 3003 3005 3007 3009 3011 3013 3015 3017 3019 3021 3023 3025 3027 3029 3031 3033 3035 3037 3039 3041 3043 3045 3047 3049 3051 3053 3055 3057 3059 3061 3063 3065 3067 3069 3071 3073 3075 3077 3079 3081 3083 3085 3087 3089 3091 3093 3095 3097 3099 3101 3103 3105 3107 3109 3111 3113 3115 3117 3119 3121 3123 3125 3127 3129 3131 3133 3135 3137 3139 3141 3143 3145 3147 3149 3151 3153 3155 3157 3159 3161 3163 3165 3167 3169 3171 3173 3175 3177 3179 3181 3183 3185 3187 3189 3191 3193 3195 3197 3199 3201 3203 3205 3207 3209 3211 3213 3215 3217 3219 3221 3223 3225 3227 3229 3231 3233 3235 3237 3239 3241 3243 3245 3247 3249 3251 3253 3255 3257 3259 3261 3263 3265 3267 3269 3271 3273 3275 3277 3279 3281 3283 3285 3287 3289 3291 3293 3295 3297 3299 3301 3303 3305 3307 3309 3311 3313 3315 3317 3319 3321 3323 3325 3327 3329 3331 3333 3335 3337 3339 3341 3343 3345 3347 3349 3351 3353 3355 3357 3359 3361 3363 3365 3367 3369 3371 3373 3375 3377 3379 3381 3383 3385 3387 3389 3391 3393 3395 3397 3399 3401 3403 3405 3407 3409 3411 3413 3415 3417 3419 3421 3423 3425 3427 3429 3431 3433 3435 3437 3439 3441 3443 3445 3447 3449 3451 3453 3455 3457 3459 3461 3463 3465 3467 3469 3471 3473 3475 3477 3479 3481 3483 3485 3487 3489 3491 3493 3495 3497 3499 3501 3503 3505 3507 3509 3511 3513 3515 3517 3519 3521 3523 3525 3527 3529 3531 3533 3535 3537 3539 3541 3543 3545 3547 3549 3551 3553 3555 3557 3559 3561 3563 3565 3567 3569 3571 3573 3575 3577 3579 3581 3583 3585 3587 3589 3591 3593 3595 3597 3599 3601 3603 3605 3607 3609 3611 3613 3615 3617 3619 3621 3623 3625 3627 3629 3631 3633 3635 3637 3639 3641 3643 3645 3647 3649 3651 3653 3655 3657 3659 3661 3663 3665 3667 3669 3671 3673 3675 3677 3679 3681 3683 3685 3687 3689 3691 3693 3695 3697 3699 3701 3703 3705 3707 3709 3711 3713 3715 3717 3719 3721 3723 3725 3727 3729 3731 3733 3735 3737 3739 3741 3743 3745 3747 3749 3751 3753 3755 3757 3759 3761 3763 3765 3767 3769 3771 3773 3775 3777 3779 3781 3783 3785 3787 3789 3791 3793 3795 3797 3799 3801 3803 3805 3807 3809 3811 3813 3815 3817 3819 3821 3823 3825 3827 3829 3831 3833 3835 3837 3839 3841 3843 3845 3847 3849 3851 3853 3855 3857 3859 3861 3863 3865 3867 3869 3871 3873 3875 3877 3879 3881 3883 3885 3887 3889 3891 3893 3895 3897 3899 3901 3903 3905 3907 3909 3911 3913 3915 3917 3919 3921 3923 3925 3927 3929 3931 3933 3935 3937 3939 3941 3943 3945 3947 3949 3951 3953 3955 3957 3959 3961 3963 3965 3967 3969 3971 3973 3975 3977 3979 3981 3983 3985 3987 3989 3991 3993 3995 3997 3999 4001 4003 4005 4007 4009 4011 4013 4015 4017 4019 4021 4023 4025 4027 4029 4031 4033 4035 4037 4039 4041 4043 4045 4047 4049 4051 4053 4055 4057 4059 4061 4063 4065 4067 4069 4071 4073 4075 4077 4079 4081 4083 4085 4087 4089 4091 4093 4095 4097 4099 4101 4103 4105 4107 4109 4111 4113 4115 4117 4119 4121 4123 4125 4127 4129 4131 4133 4135 4137 4139 4141 4143 4145 4147 4149 4151 4153 4155 4157 4159 4161 4163 4165 4167 4169 4171 4173 4175 4177 4179 4181 4183 4185 4187 4189 4191 4193 4195 4197 4199 4201 4203 4205 4207 4209 4211 4213 4215 4217 4219 4221 4223 4225 4227 4229 4231 4233 4235 4237 4239 4241 4243 4245 4247 4249 4251 4253 4255 4257 4259 4261 4263 4265 4267 4269 4271 4273 4275 4277 4279 4281 4283 4285 4287 4289 4291 4293 4295 4297 4299 4301 4303 4305 4307 4309 4311 4313 4315 4317 4319 4321 4323 4325 4327 4329 4331 4333 4335 4337 4339 4341 4343 4345 4347 4349 4351 4353 4355 4357 4359 4361 4363 4365 4367 4369 4371 4373 4375 4377 4379 4381 4383 4385 4387 4389 4391 4393 4395 4397 4399 4401 4403 4405 4407 4409 4411 4413 4415 4417 4419 4421 4423 4425 4427 4429 4431 4433 4435 4437 4439 4441 4443 4445 4447 4449 4451 4453 4455 4457 4459 4461 4463 4465 4467 4469 4471 4473 4475 4477 4479 4481 4483 4485 4487 4489 4491 4493 4495 4497 4499 4501 4503 4505 4507 4509 4511 4513 4515 4517 4519 4521 4523 4525 4527 4529 4531 4533 4535 4537 4539 4541 4543 4545 4547 4549 4551 4553 4555 4557 4559 4561 4563 4565 4567 4569 4571 4573 4575 4577 4579 4581 4583 4585 4587 4589 4591 4593 4595 4597 4599 4601 4603 4605 4607 4609 4611 4613 4615 4617 4619 4621 4623 4625 4627 4629 4631 4633 4635 4637 4639 4641 4643 4645 4647 4649 4651 4653 4655 4657 4659 4661 4663 4665 4667 4669 4671 4673 4675 4677 4679 4681 4683 4685 4687 4689 4691 4693 4695 4697 4699 4701 4703 4705 4707 4709 4711 4713 4715 4717 4719 4721 4723 4725 4727 4729 4731 4733 4735 4737 4739 4741 4743 4745 4747 4749 4751 4753 4755 4757 4759 4761 4763 4765 4767 4769 4771 4773 4775 4777 4779 4781 4783 4785 4787 4789 4791 4793 4795 4797 4799 4801 4803 4805 4807 4809 4811 4813 4815 4817 4819 4821 4823 4825 4827 4829 4831 4833 4835 4837 4839 4841 4843 4845 4847 4849 4851 4853 4855 4857 4859 4861 4863 4865 4867 4869 4871 4873 4875 4877 4879 4881 4883 4885 4887 4889 4891 4893 4895 4897 4899 4901 4903 4905 4907 4909 4911 4913 4915 4917 4919 4921 4923 4925 4927 4929 4931 4933 4935 4937 4939 4941 4943 4945 4947 4949 4951 4953 4955 4957 4959 4961 4963 4965 4967 4969 4971 4973 4975 4977 4979 4981 4983 4985 4987 4989 4991 4993 4995 4997 4999 5001 5003 5005 5007 5009 5011 5013 5015 5017 5019 5021 5023 5025 5027 5029 5031 5033 5035 5037 5039 5041 5043 5045 5047 5049 5051 5053 5055 5057 5059 5061 5063 5065 5067 5069 5071 5073 5075 5077 5079 5081 5083 5085 5087 5089 5091 5093 5095 5097 5099 5101 5103 5105 5107 5109 5111 5113 5115 5117 5119

8/45

RESTRICTION MAP OF 5'-FLANKING  
REGION OF ~~HPV~~ SI-CASEIN GENE



1	104X	X
2	105I	U
3	106N	N
4	107M	M
5	108H	H
6	109R	R
7	110G	G
8	111S	S

6

9/45

restriction map of 3'-flanking region of  $\alpha 51$ -casein gene




R -  $\Sigma$ 012  
 BS - BS+EII  
 B - B000HII  
 H - HindIII  
 K - KpnII

F, - 2.1

146

Figure - 5A




10/45





12/45



13/45

NAME PMH-1






Fig 6

Fig 7A

- Plasmid **PMH3'E10** contains a 8.5kb **EcoRI** fragment of the 3'-END OF NS1-CASEIN. PMH-1 WAS CUT W/ **EcoRI** + LIGATED TO THIS FRAGMENT



- Plasmid **PMH3'UTRHLF2 LINKER**:




14/45

15/95

CONSTRUCTION OF PSI3' hLF<sub>1/2</sub>


Fig 7-6





17/45


Construction of  $\overline{\text{PLF } 3' \text{ } 10 \text{ kb}}$




18/45

CONSTRUCTION OF PSI 3'S'HLF | Fig 7-E

STARTED w/ PSI 3'UTRHLF CUT w/ HINDIII AND SALI + PURIFIED THIS  
FRAGMENT CONTAINING THE PSI-CASEIN SIGNAL SEQUENCE + hLF CODING REGION, + 3'S'UTR  
AND SW(A)U



SUBCLONED INTO HINDIII/SALI  
CUT PMH 3'S'FLANK TO YIELD :




NOTE: PSI 3'S'HLF WAS MADE BY CUTTING PSI 3'S'HLF w/ KpnI AND BamHI,  
FOLLOWED BY BLUNTED THE ENDS AND RELIGATING. THIS ELIMINATES THE  
SPICED 3'UTR REGION





19/45

cosmid "GP1 HLF" 15, 7-F




3 way ligation. The DNA from this cosmid is prepared by cutting with **NotI** and purifying the expression sequence prior to microinjection

20/45



F<sub>18</sub> - 8



S=SalI R=NotI Bg=BglII N=NotI X=XhoI

Fig 9

22/45

phage containing protein C gene:




Fig 10

23/45

Fig-11

Shown below is the DNA sequence of the hybrid splice. The predicted intervening sequence (shown in lower case) consists of the 5'-end of IVS-1 from bovine  $\alpha$ 1 casein (from position +54 to +180 with respect to the start of transcription) fused to the 3'-end of a human IgG splice sequence. The Hind III site (in bold type and underlined) derives from the IgG sequence and marks the junction between the  $\alpha$ 1 and IgG splice sequences. The 5'-end upper case sequence depicts the complete exon one of the bovine  $\alpha$ 1 casein gene. The 3'-end upper case sequence represents the splice junction of the IgG gene through to the Pst I site (CTGCAG) incorporated in the cloning vector, pMHI.

5'- ATCACCTTGA TCATCAACCC AGCTTGCTGC TTCTTCCCAG  
 TCTTGGGTTC AAG gtattatgta tacatataac aaaatttcta tgattttct cgtctctc  
 ttcatctt cactaatagc cagttgtaac ttctatgt gatgcaagt atgggtactt tctatgata  
 tactgttagc aagcttgagg tgtggcggc ttgagatctg gccatacact tgagtgacaa tgacatccac  
 ttgccttc tctccacg GTGTCCACTC CCAGGTCCAA CTGCAG - 3'

24/45




FIG. 12A




FIG. 12B

25/45




FIG. 13

26/45




FIG. 14



27/45




FIG. 15A

28/45

Structure of Cla I - Apa I Synthetic sequence

Translation codon.

Cla I


5'-CGA TAC CAA GTC GCC TCC AGA CCG CAG ACA TGA AAC TTG TCT  
 T ATG GTT CAG CGG AGG TCT GCC GTC TGT ACT TTG AAC AGA

TCC TCG TCC TGC TGT TCC TCG GGG CC-3'  
 AGG AGC AGG ACG ACA AGG AGC C

Apa I

FIG. 15B

29/45



Structure of region containing exon 1 (hybrid 9S-cas/low. 9S-cas) and part of exon 2 of the genomic low. 9S-cas depicted in Figs. 15A through 17.

FIG. 15C

30/45




FIG. 16

31/45


Generation of  $\delta$ hLF gene

FIG. 17

32/45




FIG. 18




FIG. 19

34/45




FIG. 20



35/45





FIG. 21

36/45

Design of 16,8 hL23

Legend: see figure 9.



(Signal sequence: hL2)

FIG. 22




FIG. 23A

38/45




FIG. 23B

39/45

linker S<sub>1</sub>/S<sub>2</sub>:Sal I<sup>⊗</sup> - Not I - Sal I<sup>⊗</sup>

⊗ : = destroyed site.






FIG. 23C

40/45

Cut p0.7 8h12 Sal I



↓ ligate S1-S2




FIG. 23D

41/45




FIG. 23E

[illegible]

↓  
translation initiation codon.

FIG. 24



43/45

Fig. 25  
 Embodiment of 278/279

start site (95, signal sequence)

translation

Clal I

CGATTAACCATGAAACCTTCTATCCTCACCTGTCTGTGCTGCTCTTGA

TATT → etc

95, 1 → 912 sequence

Bcl I


CCAAAGGTCCTTGAAGGCTGTGAGTTGC

etc, → AACC

44/45


Construction of 16.8 A hLZ 3:

FIG 26



Construction of 16. A hLZ 3:

Fig. 27



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/05724

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07H 21/00; C12N 15/00; A23C 9/00; A01N 1/02.

US CL : 536/27; 435/320.1; 935/22

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/27; 435/172.3, 320.1, 2; 800/2; 426/580; 935/22

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	J. Cell Biochem., Volume Suppl. 0 (10 Part A), Issued 1986, Chisari et al., "Hepatitis B virus gene expression in transgenic mice", page 212, see the entire abstract.	<u>1</u> 2, 4-6
<u>X</u> Y	J. Cell Biol., Volume 107 (6 Part 3), Issued 1988, Steinhilper et al., "Expression of albumin-ANF and albumin-ANF receptor fusion genes in transgenic mice", page 481A, see the entire abstract.	<u>1</u> 2, 4-6

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search	Date of mailing of the international search report
03 August 1993	01 SEP 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer JASEMINE C. CHAMBERS
Facsimile No. NOT APPLICABLE	Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

Int. l. application No.  
PCT/US93/05724

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	Genes & Development, Volume 1, Issued 1987, Pinket et al., "An albumin enhancer located 10Kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice", pages 268-276, see the entire document.	<u>1</u> 2, 4-6
<u>X</u> Y	Genes & Development, Volume 1, Issued 1987, Kelsey et al., "Species-and tissue-specific expression of human $\alpha$ 1-antitrypsin in transgenic mice", pages 161-171, see the entire document.	<u>1</u> 2, 4-6
<u>X</u> Y	Genes & Development, Volume 2, Issued 1988, Dente et al., "Expression of human $\alpha$ 1-acid glycoprotein genes in cultured cells and in transgenic mice", pages 259-266, see the entire document.	<u>1</u> 2, 4-6
<u>X</u> Y	US,A, 4,873,316, (Meade et al.), 10 October 1989, see the entire document.	1, 3-6, 9, <u>10,14,18-23</u> 7,8,11-13,15- 17,24-28,63- 72,75-81
Y	J. Biol. Chem, Volume 261, No. 15, Issued 25 May 1986, Minghetti et al., "Molecular structure of the human albumin gene is revealed by nucleotide sequence within q11-22 of chromosome 4", pages 6747-6757, see the entire document.	2,4-6,15,16
Y	EP,A, 0,264,166 (Gordon et al.), 20 April 1988, see the entire document.	1-28,63-72,75-81
Y	WO,A, 88/01648 (Hopp), 10 March 1988, see the entire document.	1-28,63-72,75-81
Y	J. Biol. Chem., Volume 261, No. 7, Issued 05 March 1986, Urano et al., "The human albumin gene", pages 3244-3251, see the entire document.	2,4-6,15,16
Y	Bio/Technology, Volume 5, Issued November 1987, Gordon et al., "Production of human tissue plasminogen activator in transgenic mouse milk", pages 1183-1187, see the entire document.	1-28,63-72,75-81
Y	Bio/Technology, Volume 6, Issued February 1988, Simons et al., "Gene transfer in sheep", pages 179-183, see the entire document.	1-28,63-72,75-81
Y	Bio/Technology, Volume 7, Issued May 1989, Clark et al., "Expression of human anti-hemophilic factor IX in the milk of transgenic sheep", pages 487-492, see the entire document.	1-28,63-72,75-81

## INTERNATIONAL SEARCH REPORT

### h. National application No.

PCT/US93/05724

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. Dairy Sci., Volume 72, Issued 1989, Bremel et al., "Alteration of milk composition using molecular genetics", pages 2826-2833, see the entire document.	1-28,63-72,75-81
Y	Theriogenology, Volume 25, No. 1, Issued January 1986, Loskutoff et al., "Gene microinjection of bovine embryos facilitated by centrifugation", page 168, see the entire abstract.	1-28,63-72,75-81
Y	Theriogenology, Volume 29, No. 1, issued January 1988, Biery et al., "Gene transfer by pronuclear injection in the bovine", page 224, see the entire abstract.	1-28,63-72,75-81

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/05724

### B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

DIALOG (Files 154, 55, 311, 312)

U.S. Automated Patent System (File USPAT, 1975-1993).

Search terms: transgene, transgenic, cow, milk, bovine, albumin, casein, lactoferrin, mammary, liver, secretory, inventors' names

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claims 1-28, 63-72 and 75-81, drawn to a transgene, classified in Class 536, subclasses 27, for example.

Group II, claims 29-40, 54, 82 and 83, drawn to a transgenic bovine species and a method of making the same, classified in Classes 800 and 435, subclasses 2 and 172.3 respectively, for example.

Group III, claims 41-53, drawn to milk from a transgenic bovine species, classified in Class 426, subclass 580, for example.

Group IV, claims 55-62, drawn to a method of producing a transgenic non-human mammal having a desired phenotype, classified in Class 800, subclass 2, for example.

Group V, claim 73, drawn to a transgenic bovine species, classified in Class 800, subclass 2, for example.

Group IV, claim 74, drawn to semen of a transgenic bovine, classified in Class 435, subclass 2, for example.

Groups I, II, III, V and VI are each directed to a different and independent product. Group IV is directed to a method of producing a transgenic non-human mammal with a desired phenotype. The method of Group V is not required to produce the transgenic bovine of either Group II or V. Accordingly, the claims are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept. See PCT Rule 13.2(i).